

# A non-conservative Harris ergodic theorem

Vincent Bansaye      Bertrand Cloez      Pierre Gabriel  
Aline Marguet

## Abstract

We consider non-conservative positive semigroups and obtain necessary and sufficient conditions for uniform exponential contraction in weighted total variation norm. This ensures the existence of Perron eigenelements and provides quantitative estimates of spectral gaps, complementing Krein-Rutman theorems and generalizing probabilistic approaches. The proof is based on a non-homogenous  $h$ -transform of the semigroup and the construction of Lyapunov functions for this latter. It exploits then the classical necessary and sufficient conditions of Harris's theorem for conservative semigroups and recent techniques developed for the study for absorbed Markov process. We apply these results to population dynamics. We obtain exponential convergence of birth and death processes conditioned on survival to their quasi-stationary distribution, as well as estimates on exponential relaxation to stationary profiles in growth-fragmentation PDEs.

**Keywords.** Positive semigroups; measure solutions; ergodicity; Krein-Rutman theorem; branching processes; growth-fragmentation; quasi-stationary distribution

## Contents

### 1 Introduction

2

---

F1. Bansaye (corresponding author): CMAP, École Polytechnique, Route de Saclay, 91128 Palaiseau cedex, France; e-mail: [vincent.bansaye@polytechnique.edu](mailto:vincent.bansaye@polytechnique.edu)

F2. Cloez: MISTEA, Univ Montpellier, INRAE, Institut Agro, Montpellier, France; e-mail: [bertrand.cloez@inra.fr](mailto:bertrand.cloez@inra.fr)

F3. Gabriel: Laboratoire de Mathématiques de Versailles, UVSQ, CNRS, Université Paris-Saclay, 45 Avenue des États-Unis, 78035 Versailles cedex, France; e-mail: [pierre.gabriel@uvsq.fr](mailto:pierre.gabriel@uvsq.fr)

F4. Marguet: Univ. Grenoble Alpes, Inria, 38000 Grenoble, France; e-mail: [aline.marguet@inria.fr](mailto:aline.marguet@inria.fr)

*Mathematics Subject Classification (2010):* Primary 47A35; Secondary 35B40, 47D06, 60J80, 92D25

<b>2</b>	<b>Definitions and main result</b>	<b>8</b>
2.1	Assumptions and criterion for exponential convergence . . . . .	9
2.2	Sufficient conditions: drift and irreducibility . . . . .	10
<b>3</b>	<b>Quantitative estimates</b>	<b>12</b>
3.1	The embedded conservative semigroup . . . . .	12
3.2	Estimation of the eigenelements . . . . .	14
3.3	Uniform exponential convergence . . . . .	15
<b>4</b>	<b>Proofs</b>	<b>15</b>
4.1	Preliminary inequalities . . . . .	15
4.2	Contraction property: proofs of Section 3.1 . . . . .	17
4.3	Eigenelements: proofs of Section 3.2 . . . . .	20
4.4	Proofs of Section 3.3 . . . . .	27
4.5	Proofs of Section 2 . . . . .	29
<b>5</b>	<b>Applications</b>	<b>33</b>
5.1	Convergence to quasi-stationary distribution . . . . .	33
5.2	The growth-fragmentation equation . . . . .	36
5.3	Comments and a few perspectives . . . . .	43
<b>6</b>	<b>Appendix</b>	<b>45</b>
6.1	Conservative operators . . . . .	45
6.2	Localization argument . . . . .	47
6.3	The growth-fragmentation semigroup . . . . .	47

# 1 Introduction

Iteration of a positive linear operator is a fundamental issue in operator analysis, linear Partial Differential Equations (PDEs), probabilities, optimization and control. In continuous time, the associated structure is a positive semigroup. A positive semigroup is a family of linear operators  $(M_t)_{t \in \mathbb{R}_+}$  acting on a space of measurable functions  $f$  on  $\mathcal{X}$ . It satisfies  $M_0 f = f$ ,  $M_{t+s} f = M_t(M_s(f))$  for any  $t, s \geq 0$ , and  $M_t f \geq 0$  for any  $f \geq 0$  and  $t \geq 0$ . In finite dimension, *i.e.* when  $\mathcal{X}$  is a finite set, under a strong positivity assumption, the Perron-Frobenius theorem [71, 43] ensures the existence of unique right and left positive eigenvectors  $h = (h(x) : x \in \mathcal{X})$  and  $\gamma = (\gamma(x) : x \in \mathcal{X})$  associated to the maximal eigenvalue  $\lambda \in \mathbb{R} : M_t h = e^{\lambda t} h$  and  $\gamma M_t = e^{\lambda t} \gamma$ . Moreover, the asymptotic profile when  $t \rightarrow +\infty$  is given by

$$M_t f = e^{\lambda t} \langle \gamma, f \rangle h + \mathcal{O}(e^{(\lambda - \omega)t}),$$

for any vector  $f$ , any  $\omega > 0$  strictly smaller than the spectral gap and  $\langle \gamma, f \rangle = \sum_{x \in \mathcal{X}} \gamma(x) f(x)$ .

The generalization in infinite dimension has attracted lots of attention. It is motivated in particular by the asymptotic analysis of linear PDEs counting the density of

particules (or individuals)  $u_t^x(y)$  in location  $y$  at time  $t$  when initially the particles are located in  $x$ , *i.e.*  $u_0^x = \delta_x$ . The semigroup is then given by  $M_t f(x) = \int_{\mathcal{X}} u_t^x(y) f(y) dy$ . Semigroups play also a key role in the study of the convergence in law of Markov processes via  $M_t f(x) = \mathbb{E}_x[f(X_t)]$  or the study of quasi-stationary regimes via  $M_t f(x) = \mathbb{E}_x[f(X_t) \mathbf{1}_{X_t \notin \mathcal{D}}]$ , where  $\mathcal{D}$  is an absorbing domain. In spectral theory, the Krein-Rutman theorem [57] yields an extension to positive compact operators for the existence of eigenelements. Various generalizations have been then obtained [51, 69, 27], the most recent providing asymptotic profiles with exponential convergence associated to a spectral gap, see in particular [68] and the references therein. When positive eigenelements exist, an alternative tool to prove convergence is the dissipation of entropy, especially the general relative entropy introduced in [67].

Our aim is to relax and simplify some assumptions involved in the spectral approach and provide more quantitative results. We apply them to two classical models: growth fragmentation PDEs and birth and death processes.

For that purpose, we are inspired and use techniques developed in probabilities. When  $M_t \mathbf{1} = \mathbf{1}$  for any  $t \geq 0$ , the semigroup is said to be *conservative*. Indeed, the total number of particles is preserved along time for the corresponding linear PDE or for the first moment semigroup of the branching Markov process. It holds when particles move without reproduction or death. Conservative semigroups arise classically in the study of the law of Markov processes  $(X_t)_{t \geq 0}$  via  $M_t f(x) = \mathbb{E}_x[f(X_t)]$ . In that case, the ergodic theory and stability of Markov processes can be invoked to analyze the long time behavior of the semigroup, in the spirit of the pioneering works of Doeblin [33] and Harris [50] and coupling techniques. We refer to [65] for an overview. These results allow proving the existence of a stationary probability measure and the uniform exponential convergence for the weighted total variation distance towards this latter. They rely on the two following conditions, corresponding respectively to Lyapunov function and small set property, for some time  $\tau > 0$  and some subset  $K$  of  $\mathcal{X}$ . First, there exists  $V : \mathcal{X} \rightarrow [1, +\infty)$  such that

$$M_\tau V \leq \mathbf{a}V + \mathbf{b}\mathbf{1}_K \quad (1.1)$$

for some constants  $\mathbf{a} \in (0, 1)$  and  $\mathbf{b} \geq 0$ . Furthermore, there exists a probability measure  $\nu$  on  $\mathcal{X}$  such that for any non-negative  $f$  and any  $x \in K$ ,

$$M_\tau f(x) \geq \mathbf{c}\nu(f) \quad (1.2)$$

with  $\mathbf{c} > 0$ . Under aperiodicity condition, these assumptions ensure that  $M_\tau$  admits a unique invariant probability measure  $\gamma$  and satisfies a contraction principle for the weighted total variation distance, see [65, 66, 56]. In our study, we will assume that

the probability measure  $\nu$  is supported by the small set  $K$ , thus guaranteeing aperiodicity. Under mild conditions of local boundedness, the contraction property captures the continuous time by iteration. Then there exist explicit constants  $C, \omega > 0$  such that for all  $x \in \mathcal{X}$ ,  $f$  such that  $|f| \leq V$ , and all  $t \geq 0$ ,

$$|M_t f(x) - \gamma(f)| \leq CV(x)e^{-\omega t},$$

where  $\gamma(f) = \int_{\mathcal{X}} f d\gamma$ . It provides a necessary and sufficient condition and quantitative estimates [36, Chapter 15] and is referred to as  $V$ -uniform ergodicity [65].

**Main results and ideas of the proof.** In the present work we provide a counterpart to  $V$ -uniform ergodicity in the non-conservative framework. We obtain necessary and sufficient conditions for exponential convergence. We also estimate the constants involved, in particular the amplitude of the spectral gap, using only the parameters in the assumptions. The approach relies on the reduction of the problem to a non-homogeneous but conservative one.

In this vein, the Doob  $h$ -transform is a historical and inspiring technique [34, 69, 35, 74]. In the case when we already know some positive eigenfunction  $h$  associated with the eigenvalue  $\lambda$ , a conservative semigroup is indeed defined for  $t \geq 0$  by

$$P_t f = \frac{M_t(hf)}{e^{\lambda t} h}.$$

The ergodic analysis of the conservative semigroup  $P$  using the theory mentioned above yields the expected estimates on the original semigroup  $M_t f = e^{\lambda t} h P_t(f/h)$ .

Our work focuses on the case when we do not know *a priori* the existence of positive eigenfunctions. We simultaneously prove their existence and estimate them. The novelty of the approach mainly lies in the form of the  $h$ -transform and the general and explicit construction of Lyapunov functions for this latter. It also strongly exploits recent progresses on the convergence to quasistationary distribution. More precisely, we introduce the transform for  $s \leq u \leq t$ :

$$P_{s,u}^{(t)} f = \frac{M_{u-s}(f M_{t-u} \psi)}{M_{t-s} \psi}.$$

For a fixed time  $t$  and positive function  $\psi$ , this family is a non-homogeneous conservative semigroup. To mimic the stabilizing property of the eigenfunction  $h$  in the  $h$ -transform, we assume that  $\psi$  satisfies

$$M_\tau \psi \geq \beta \psi, \tag{1.3}$$

for some  $\beta > 0$  and  $\tau > 0$ , similarly as in [25, 68, 22]. Section 5 provides examples where  $\psi$  can be obtained, sometimes explicitly, while we do not know a priori

eigenelements. We construct a family of Lyapunov functions  $(V_k)_{k \geq 0}$  as follows

$$V_k = \nu \left( \frac{M_{k\tau}\psi}{\psi} \right) \frac{V}{M_{k\tau}\psi},$$

where  $\nu$  is a probability measure. Assuming that  $V$  satisfies

$$M_\tau V \leq \alpha V + \theta \mathbf{1}_K \psi, \tag{1.4}$$

these functions allow us to prove that  $P^{(t)}$  satisfies Lyapunov type condition (1.1). To prove that  $P^{(t)}$  also satisfies the small set condition (1.2) and invoke the contraction of the conservative framework, we follow techniques due to Champagnat and Villemonais [21] for the study for the convergence of processes conditioned on non-absorption. This yields the following generalized small set condition. We assume that there exist  $c, d > 0$  such that for all  $x \in K$ ,

$$M_\tau(f\psi)(x) \geq c \nu(f) M_\tau\psi(x) \tag{1.5}$$

for any  $f$  positive and

$$M_{n\tau}\psi(x) \leq \frac{1}{d} \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) \psi(x) \tag{1.6}$$

for any integer  $n$ . The two latter conditions involve uniform but local bounds on the space  $\mathcal{X}$ . The first inequality is restricted to a fixed time. Convenient sufficient condition can be given to restrict also the second one to a finite time estimate, in particular via coupling, see the next paragraph and applications.

Assuming that  $\alpha = \alpha/\beta < 1$  and  $V \geq \psi$ , we obtain a contraction result for  $P^{(t)}$  in forthcoming Proposition 3.3. By estimating the mass  $M_t\psi$ , we then prove that a triplet  $(\lambda, h, \gamma)$  of eigenelements exists and provide estimates on these elements. We get that for all  $x \in \mathcal{X}$  and  $f$  such that  $|f| \leq V$ ,

$$|e^{-\lambda t} M_t f(x) - h(x)\gamma(f)| \leq CV(x)e^{-\omega t},$$

with constants  $\omega > 0$  and  $C > 0$ , that are explicit. We also prove that the conditions given above are actually necessary for this uniform convergence in weighted total variation distance.

**Tractable conditions and applications.** We apply our results to describe the profile of populations in growth-fragmentation PDEs and birth and death processes conditioned on non-absorption. We also refer to the follow-up paper [24] for an application to mutation-selection PDEs.

Sufficient conditions are helpful for such issues. In particular, one can check Lyapunov conditions (1.3) and (1.4) using the generator. Loosely speaking, writing  $\mathcal{L}$  the generator of  $(M_t)_{t \geq 0}$ , we check that it is enough to find  $V, \psi$  and  $a < b$  such that

$$\mathcal{L}V \leq aV + \zeta\psi \quad \text{and} \quad \mathcal{L}\psi \geq b\psi. \quad (1.7)$$

The second part of the assumptions deals with local bounds on the semigroup. It is worth noticing that in the case  $0 < \inf_K \psi \leq \sup_K \psi < \infty$ , Assumption (1.5) is equivalent to the small set condition (1.2). For a discrete state space  $\mathcal{X}$ , the conditions (1.5) and (1.6) can be reduced to irreducibility properties. In the continuous setting, a coupling condition implying (1.6) is proposed and applied in [24]:

$$\frac{M_\tau f(x)}{\psi(x)} \geq c_0 \int_0^\tau \frac{M_{\tau-s} f(y)}{\psi(y)} \sigma_{x,y}(ds)$$

for all  $x, y \in K$ ,  $f \geq 0$  and some constants  $c_0, \tau > 0$  and probability measures  $(\sigma_{x,y})_{x,y \in K}$  on  $[0, \tau]$ . This condition is stronger than Assumption (1.6), but it is easier to check for several population models.

Let us briefly illustrate the interest of the conditions above and the novelty of the result for applications on the so-called growth-fragmentation equation. This equation is central in the modeling of various physical or biological phenomena [37, 64, 72, 3, 75, 4]. In its classical form it can be written as

$$\partial_t u_t(x) + \partial_x u_t(x) + B(x)u_t(x) = \int_0^1 B\left(\frac{x}{z}\right) u_t\left(\frac{x}{z}\right) \frac{\wp(dz)}{z},$$

where  $t, x > 0$ , and is complemented with a zero flux boundary condition  $u_t(0) = 0$ . It drives the time evolution of the population density  $u_t$  of particles characterized by a structural size  $x \geq 0$ . Each particle with size  $x$  grows with speed one and splits with rate  $B(x)$  to produce smaller particles of sizes  $zx$ , with  $0 < z < 1$  distributed with respect to the fragmentation kernel  $\wp$ . The corresponding dual equation reads

$$\partial_t \varphi_t(x) = \mathcal{L} \varphi_t(x), \quad \text{where} \quad \mathcal{L} f(x) = f'(x) + B(x) \left( \int_0^1 f(zx) \wp(dz) - f(x) \right).$$

It generates a semigroup  $(M_t)_{t \geq 0}$  on the state space  $\mathcal{X} = [0, \infty)$  as follows:  $M_t f = \varphi_t$  is the solution to the dual equation with initial condition  $\varphi_0 = f$ . The duality property ensures the relation  $\int_{\mathcal{X}} u_t(x) f(x) dx = \int_{\mathcal{X}} u_0(x) M_t f(x) dx$  for all time  $t \geq 0$ . A direct computation ensures that the infinitesimal generator  $\mathcal{L}$  verifies (1.7) with

$$V : x \mapsto 1 + x^k, \quad k > 1 \quad \text{and} \quad \psi : x \mapsto 1 + x.$$

Consequently the semigroup  $(M_t)_{t \geq 0}$  satisfies (1.3)-(1.4). Using the reachability of any sufficiently large size and a monotonicity argument on  $B$ , we can show (1.5)-(1.6). It gives the existence of eigenelements and the exponential convergence. It improves the existing results we know, where a polynomial bound is assumed for  $B$ . It additionally provides estimates on the spectral gap. We refer the reader to Section 5.2 for details and more references.

**State of the art and related works.** Our results and method are linked to the study of geometric ergodicity of Feynman-Kac type semigroups

$$M_t f(x) = \mathbb{E}_x \left[ f(X_t) e^{\int_0^t F(X_s) ds} \right].$$

The study of these non-conservative semigroups has been developed in a general setting [54, 55]. We refer also to [31, 30] in discrete time. These semigroups appear in particular for the study of branching processes [23, 62, 15] and large deviations [54, 55, 41]. In [54] the function  $F$  is supposed to be bounded, with a norm smaller than an explicit constant. The unbounded case is addressed in [55] where some weighted norm of  $F$  is required to be small, together with either a “density assumption” or some irreducibility and aperiodicity conditions. The growth-fragmentation model would correspond to  $F = B$  and we relax here the assumptions on  $B$ . Nevertheless, we observe that the criteria in [54, 55] may be simpler to apply, and under the density assumption the authors prove a stronger result of discrete spectrum.

In the same vein, let us mention [42], which uses the Krein-Rutman theorem for the existence of eigenelements and an  $h$ -transform. It requires additional regularity assumptions to derive the geometric ergodicity of general Feynman-Kac semigroups, as well as a stronger small set condition. The regularity requirement is a local strong Feller assumption. It is well-suited for diffusive equations, see also [53] for a result on non-conservative hypoelliptic diffusions. It does not seem to be adapted to our motivations with less regularizing effect.

In [14, 15], a Feynman-Kac approach is developed for growth-fragmentation equation with exponential individual growth. It relies on the application of the Krein-Rutman theorem on an well-suited operator. It provides sharp conditions on the growth rate but it requires the division rate  $B$  to be bounded. We expect that our results provide quantitative estimates and relax the boundedness assumption on  $B$  in [14, 15] for some relevant classes of growth rates.

Finally, Hilbert metric and Birkhoff contraction yield another powerful method for analysis of semigroup, which has been well developed [16, 76, 70]. As far as we see, it requires uniform bounds on the whole space for the semigroup at fixed time, which are relaxed by the backward approach exploited here.

More generally, our work is motivated by the study of structured populations and varying environment. The conditions are well adapted to complex trait spaces for populations, including discrete and continuous components: space, age, phenotype, genotype, size, etc., see Section 5 and *e.g.* [9, 61, 8, 24]. The contraction with explicit bounds allows relevant compositions appearing when parameters of the population dynamic vary along time. Such a method has already been exploited in [8] for the analysis of PDEs in varying environment in the case of uniform exponential convergence. Several models need the more general framework considered in this paper. Indeed, the typical trait does not come back in compact sets in a bounded time for models like growth-fragmentation and mutation-selection.

**Outline of the paper.** We stress that the paper is self-contained. It is structured as follows. In Section 2, we gather the assumptions, the necessary and sufficient condition in weighted total variation distance and useful sufficient conditions. In Section 3.1, we consider the non-homogeneous conservative embedded semigroup and we establish its contraction property. We present in Section 3.2 our results on the existence and the estimations of the eigenelements and in Section 3.3 complementary convergence estimates. Section 4 contains the proofs of these statements. Section 5 is devoted to applications: the random walk on integers absorbed at 0 and the growth-fragmentation equation. We finally mention briefly some additional results and perspectives, including the discrete time framework and the reducible case.

## 2 Definitions and main result

We start by stating precisely our framework. Let  $\mathcal{X}$  be a measurable space. For any measurable function  $\varphi : \mathcal{X} \rightarrow (0, \infty)$  we denote by  $\mathcal{B}(\varphi)$  the space of measurable functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  which are dominated by  $\varphi$ , *i.e.* such that the quantity

$$\|f\|_{\mathcal{B}(\varphi)} = \sup_{x \in \mathcal{X}} \frac{|f(x)|}{\varphi(x)}$$

is finite. Endowed with this weighted supremum norm,  $\mathcal{B}(\varphi)$  is a Banach space. Let  $\mathcal{B}_+(\varphi) \subset \mathcal{B}(\varphi)$  be its positive cone, namely the subset of nonnegative functions.

Let  $\mathcal{M}_+(\varphi)$  be the cone of positive measures on  $\mathcal{X}$  which integrate  $\varphi$ , *i.e.* the set of positive measures  $\mu$  on  $\mathcal{X}$  such that the quantity  $\mu(\varphi) = \int_{\mathcal{X}} \varphi d\mu$  is finite. We denote by  $\mathcal{M}(\varphi) = \mathcal{M}_+(\varphi) - \mathcal{M}_+(\varphi)$  the set of the differences of measures that belong to  $\mathcal{M}_+(\varphi)$ . When  $\inf \varphi > 0$ , this set is the subset of signed measures which integrate  $\varphi$ . When  $\inf \varphi = 0$  it is not a subset of signed measures, but in any case it



can be rigorously built as the quotient space

$$\mathcal{M}(\varphi) = \mathcal{M}_+(\varphi) \times \mathcal{M}_+(\varphi) / \sim$$

where  $(\mu_1, \mu_2) \sim (\tilde{\mu}_1, \tilde{\mu}_2)$  if  $\mu_1 + \tilde{\mu}_2 = \mu_2 + \tilde{\mu}_1$ , and we denote  $\mu = \mu_1 - \mu_2$  to mean that  $\mu$  is the equivalence class of  $(\mu_1, \mu_2)$ . The Hahn-Jordan Decomposition Theorem ensures that for any  $\mu \in \mathcal{M}(\varphi)$  there exists a unique couple  $(\mu_+, \mu_-) \in \mathcal{M}_+(\varphi) \times \mathcal{M}_+(\varphi)$  of mutually singular measures such that  $\mu = \mu_+ - \mu_-$ . An element  $\mu$  of  $\mathcal{M}(\varphi)$  acts on  $\mathcal{B}(\varphi)$  through  $\mu(f) = \mu_+(f) - \mu_-(f)$  and  $\mathcal{M}(\varphi)$  is endowed with the weighted total variation norm

$$\|\mu\|_{\mathcal{M}(\varphi)} = \mu_+(\varphi) + \mu_-(\varphi) = \sup_{\|f\|_{\mathcal{B}(\varphi)} \leq 1} |\mu(f)|$$

which makes it a Banach space.

We are interested in semigroups  $(M_t)_{t \geq 0}$  of kernel operators, *i.e.* semigroups of linear operators  $M_t$  which act both on  $\mathcal{B}(\varphi)$  (on the right  $f \mapsto M_t f$ ) and on  $\mathcal{M}(\varphi)$  (on the left  $\mu \mapsto \mu M_t$ ), are positive in the sense that they leave invariant the positive cones of both spaces, and enjoy the duality relation  $(\mu M_t)(f) = \mu(M_t f)$ . We work in the continuous time setting,  $t \in \mathbb{R}^+$ , and we refer to Section 5.3 for comments on the discrete time case. Finally, we use the shorthand notation  $f \lesssim g$  when, for two functions  $f, g : \Omega \rightarrow \mathbb{R}$ , there exists a constant  $C > 0$  such that  $f \leq Cg$  on  $\Omega$ .

## 2.1 Assumptions and criterion for exponential convergence

We consider two measurable functions  $V : \mathcal{X} \rightarrow (0, \infty)$  and  $\psi : \mathcal{X} \rightarrow (0, \infty)$ , with  $\psi \leq V$ , and a positive semigroup  $(M_t)_{t \geq 0}$  of kernel operators acting on  $\mathcal{B}(V)$  and  $\mathcal{M}(V)$ . Let us state Assumption **A**, which gathers conditions given in Introduction.

**Assumption A.** There exist  $\tau > 0$ ,  $\beta > \alpha > 0$ ,  $\theta > 0$ ,  $(c, d) \in (0, 1]^2$ ,  $K \subset \mathcal{X}$  and  $\nu$  a probability measure on  $\mathcal{X}$  supported by  $K$  such that  $\sup_K V/\psi < \infty$  and

$$(A1) \quad M_\tau V \leq \alpha V + \theta \mathbf{1}_K \psi,$$

$$(A2) \quad M_\tau \psi \geq \beta \psi,$$

$$(A3) \quad \inf_{x \in K} \frac{M_\tau(f\psi)(x)}{M_\tau \psi(x)} \geq c \nu(f) \quad \text{for all } f \in \mathcal{B}_+(V/\psi),$$

$$(A4) \quad \nu \left( \frac{M_{n\tau} \psi}{\psi} \right) \geq d \sup_{x \in K} \frac{M_{n\tau} \psi(x)}{\psi(x)} \quad \text{for all positive integers } n.$$

Assumption **A** is linked and relax classical assumptions for the ergodicity of non-conservative semigroups [8, 22, 42, 53, 54, 55, 68, 79, 80].

More precisely, one can compare **A** to the assumptions (1-4) of [68, Theorem 5.3], where (2) corresponds to (A2), (1) has strong connections with (A1) (see [83]), while (4) is relaxed. Besides, the small set condition (A3) is more general than strong positivity or irreducibility assumptions.

Assumption **A** provides an extension of the conditions of [8] in the homogeneous case. In that latter, small set condition (A3) was required on the whole space  $\mathcal{X}$ , which imposes uniform ergodicity. Uniformity does not hold in the two applications we consider in the present paper.

Our assumptions are similar and inspired from [22], which is dedicated to convergence to quasi-stationary distribution. Our technique differs, especially for the form of the embedded semigroup and the construction of Lyapunov functions, see Section 3.1. We relax the boundedness of  $\psi$  required in [22] and obtain necessary conditions for weighted exponential convergence. As far as we see, this approach also provides more quantitative estimates, see forthcoming Section 3 and the proofs.

The main result of the paper can be stated as follows. It is proved in Section 4.5, where constants involved are explicit. To state the result in the continuous time setting, we need that  $M_t V(x)/V(x)$  is bounded on compact time intervals, uniformly on  $\mathcal{X}$ . We refer to [65, Section 20.3] for details on this classical assumption.

**Theorem 2.1.** *Assume that  $t \mapsto \|M_t V\|_{\mathcal{B}(V)}$  is locally bounded on  $[0, \infty)$ .*

*i) Let  $(V, \psi)$  be a couple of measurable functions from  $\mathcal{X}$  to  $(0, \infty)$  which satisfies Assumption **A**. Then, there exists a unique triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$  of eigenelements of  $M$  with  $\gamma(h) = \|h\|_{\mathcal{B}(V)} = 1$  satisfying for all  $t \geq 0$ ,*

$$\gamma M_t = e^{\lambda t} \gamma \quad \text{and} \quad M_t h = e^{\lambda t} h. \quad (2.1)$$

*Moreover, there exist  $C, \omega > 0$  such that for all  $t \geq 0$  and  $\mu \in \mathcal{M}(V)$ ,*

$$\|e^{-\lambda t} \mu M_t - \mu(h) \gamma\|_{\mathcal{M}(V)} \leq C \|\mu\|_{\mathcal{M}(V)} e^{-\omega t}. \quad (2.2)$$

*ii) Assume that there exist a positive measurable function  $V$ , a triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$ , and constants  $C, \omega > 0$  such that (2.1) and (2.2) hold. Then, the couple  $(V, h)$  satisfies Assumption **A**.*

## 2.2 Sufficient conditions: drift and irreducibility

Assumptions (A1)-(A2) can be checked more easily using the generator  $\mathcal{L}$  of the semigroup  $(M_t)_{t \geq 0}$ . We give convenient sufficient conditions by adopting a mild

formulation of  $\mathcal{L} = \partial_t M_t|_{t=0}$ , similar to [47]. For  $F, G \in \mathcal{B}(V)$  we say that  $\mathcal{L}F = G$  if for all  $x \in \mathcal{X}$  the function  $s \mapsto M_s G(x)$  is locally integrable, and for all  $t \geq 0$ ,

$$M_t F = F + \int_0^t M_s G ds.$$

In general for  $F \in \mathcal{B}(V)$ , there may not exist  $G \in \mathcal{B}(V)$  such that  $\mathcal{L}F = G$ , meaning that  $F$  is not in the domain of  $\mathcal{L}$ . Therefore we relax the definition by saying that

$$\mathcal{L}F \leq G, \quad \text{resp.} \quad \mathcal{L}F \geq G,$$

if for all  $t \geq 0$

$$M_t F - F \leq \int_0^t M_s G ds, \quad \text{resp.} \quad M_t F - F \geq \int_0^t M_s G ds.$$

We can now state the drift conditions on  $\mathcal{L}$  guaranteeing the validity of Assumptions (A1)-(A2). It will be useful for the applications in Section 5. For convenience, we use the shorthand  $\varphi \simeq \psi$  to mean that  $\psi \lesssim \varphi \lesssim \psi$ , *i.e.* that the ratios of the two functions are bounded.

**Proposition 2.2.** *Let  $V, \psi, \varphi : \mathcal{X} \rightarrow (0, \infty)$  such that  $\psi \leq V$  and  $\varphi \simeq \psi$ . Assume that there exist constants  $a < b$  and  $\zeta \geq 0, \xi \in \mathbb{R}$  such that*

$$\mathcal{L}V \leq aV + \zeta\psi, \quad \mathcal{L}\psi \geq b\psi, \quad \mathcal{L}\varphi \leq \xi\varphi.$$

*Then, for any  $\tau > 0$ , there exists  $R > 0$  such that  $(V, \psi)$  satisfies (A1)-(A2) with  $K = \{V \leq R\psi\}$ .*

We provide now a sufficient condition for (A3)-(A4).

**Proposition 2.3.** *Let  $K$  be a finite subset of  $\mathcal{X}$  and assume that there exists  $\tau > 0$  such that for any  $x, y \in K$ ,*

$$\delta_x M_\tau(\mathbf{1}_{\{y\}}) > 0.$$

*Then (A3)-(A4) are satisfied for any positive function  $\psi \in \mathcal{B}(V)$ .*

This sufficient condition is relevant for the study of irreducible processes on discrete spaces. We refer to Section 5.1 for an application to the convergence to quasi-stationary distribution of birth and death processes. As a motivation, let us also mention the study of the first moment semigroup of discrete branching processes in continuous time and the exponential of denumerable non-negative matrices. For continuous state space, the irreducibility condition above is not relevant and we refer to [24] for more general conditions to check (A4) via a coupling argument.

### 3 Quantitative estimates

To exploit the conservative theory, we consider a relevant conservative semigroup associated to  $M$ . We then derive the expected estimates for the eigenelements and obtain the quantitative estimates for the original semigroup  $M$ .

#### 3.1 The embedded conservative semigroup

Let us fix a positive function  $\psi \in \mathcal{B}(V)$  and a time  $t > 0$ . For any  $0 \leq s \leq u \leq t$ , we define the operator  $P_{s,u}^{(t)}$  acting on bounded measurable functions  $f$  through

$$P_{s,u}^{(t)}f = \frac{M_{u-s}(fM_{t-u}\psi)}{M_{t-s}\psi}. \quad (3.1)$$

We observe that the family  $P^{(t)} = (P_{s,u}^{(t)})_{0 \leq s \leq u \leq t}$  is a non-homogeneous conservative semigroup (or semiflow, or propagator), meaning that for any  $0 \leq s \leq u \leq v \leq t$ ,

$$P_{s,u}^{(t)}P_{u,v}^{(t)} = P_{s,v}^{(t)}.$$

It has a probabilistic interpretation in terms of particles systems, see *e.g.* [62] and references below. Roughly speaking, it provides the position of the backward lineage of a particle at time  $t$  sampled with a bias  $\psi$ .

The particular case  $\psi = \mathbf{1}$  corresponds to uniform sampling and has been successfully used in the study of positive semigroups, see [29, 6, 21, 62, 8, 22]. Whenever possible, the right eigenfunction of the semigroup provides a relevant choice for  $\psi$ . Indeed, if  $h$  is a positive eigenfunction, the conservative semigroup  $P^{(t)}$  associated to  $\psi = h$  is homogeneous and does not depend on  $t$ :  $P_s = P_{u,u+s}^{(t)} = M_s(h \cdot)/e^{\lambda s}h$ . This corresponds to the  $h$ -transform given in Introduction. This transformation provides a powerful tool for the analysis of branching processes and absorbed Markov process, see *e.g.* respectively [58, 39, 23] and [34, 26]. To shed some light on the sequel, let us explain how to apply Harris's ergodic theorem ([48] or Theorem 6.1 in Appendix 6.1 with  $\mathcal{W} = \mathcal{V}$ ) to  $(P_s)_{s \geq 0}$  and get the asymptotic behavior of  $M$ . Inequality (6.1) for  $P_\tau$  reads  $P_\tau \mathcal{V} \leq \alpha \mathcal{V} + \mathbf{c}$  and yields  $M_\tau(\mathcal{V}h) \leq \alpha \mathcal{V}h + \theta h$ , with  $\alpha = \mathbf{a}e^{\lambda \tau}$ ,  $\theta = \mathbf{c}e^{\lambda \tau}$ . This inequality involving  $M_\tau$  is guaranteed by (A1) by setting  $V = \mathcal{V}h$  and  $\psi = h$ . Additionally, Equation (6.2) corresponds exactly to (A3).

In this paper, we deal with the general case and consider a positive function  $\psi$  satisfying (A2). The analogy with the  $h$ -transform above suggests to look for Lyapunov functions of the form  $\mathcal{V} = V/\psi$ . The family of functions  $(V/M_{k\tau}\psi)_{k \geq 0}$  satisfies, under Assumption (A1), an extended version of the Lyapunov condition

for  $P^{(t)}$ . But their level sets may degenerate as  $k$  goes to infinity, which raises a problem to check the small set condition. We compensate the magnitude of  $M_{k\tau}\psi$  and consider for  $k \geq 0$ ,

$$V_k = \nu \left( \frac{M_{k\tau}\psi}{\psi} \right) \frac{V}{M_{k\tau}\psi}. \quad (3.2)$$

The two following lemmas, which are proved in Section 4.2, ensure that  $(V_k)_{k \geq 0}$  provides Lyapunov functions whose sublevel sets are small for  $P^{(t)}$ .

**Lemma 3.1.** *For all integers  $k \geq 0$  and  $n \geq m \geq k + 1$ , we have*

$$P_{k\tau, m\tau}^{(n\tau)} V_{n-m} \leq \mathbf{a} V_{n-k} + \mathbf{c},$$

where

$$\mathbf{a} = \frac{\alpha}{\beta} \in (0, 1), \quad \mathbf{c} = \frac{\theta}{c(\beta - \alpha)} \geq 0. \quad (3.3)$$

**Lemma 3.2.** *Let  $\mathfrak{R} > 0$  and set*

$$\mathbf{p} = \left\lfloor \frac{\log \left( \frac{2\mathfrak{R}(\alpha + \theta)}{(\beta - \alpha)d} \right)}{\log(\beta/\alpha)} \right\rfloor + 1. \quad (3.4)$$

Then, there exists a family of probability measures  $\{\nu_{k,n}, k \leq n\}$  such that for all  $0 \leq k \leq n - \mathbf{p}$  and  $x \in \{V_{n-k} \leq \mathfrak{R}\}$ ,

$$\delta_x P_{k\tau, (k+\mathbf{p})\tau}^{(n\tau)} \geq \mathbf{b} \nu_{k,n},$$

where  $\mathbf{b} \in (0, 1]$  is defined by

$$\mathbf{b} = \frac{d^2 \beta}{2c^2(\alpha/\theta + 1)(\alpha R + \theta)} \frac{1}{\sum_{j=1}^{\ell} (\mathbf{a}/cr)^j},$$

with  $R = \sup_K V/\psi$  and  $r = (\beta/(\alpha(R + \theta/(\beta - \alpha)) + \theta))^2$ .

We can now state the key contraction result. Its proof lies in the two previous lemmas and a slight adaptation of Harris theorem provided in Appendix.

**Proposition 3.3.** *Let  $(V, \psi)$  be a couple of measurable functions from  $\mathcal{X}$  to  $(0, \infty)$  satisfying Assumption **A**. Let  $\mathfrak{R} > 2c/(1 - \mathbf{a})$ ,  $\mathbf{b}' \in (0, \mathbf{b})$ ,  $\mathbf{a}' \in (\mathbf{a} + 2c/\mathfrak{R}, 1)$  and set*

$$\kappa = \frac{\mathbf{b}'}{c}, \quad \eta = \max \left\{ 1 - (\mathbf{b} - \mathbf{b}'), \frac{2 + \kappa \mathfrak{R} \mathbf{a}'}{2 + \kappa \mathfrak{R}} \right\}.$$

Then, for any  $\mu_1, \mu_2 \in \mathcal{M}_+(V/\psi)$  and any integers  $k$  and  $n$  such that  $0 \leq k \leq n - \mathbf{p}$ ,

$$\left\| \mu_1 P_{k\tau, (k+\mathbf{p})\tau}^{(n\tau)} - \mu_2 P_{k\tau, (k+\mathbf{p})\tau}^{(n\tau)} \right\|_{\mathcal{M}(1+\kappa V_{n-k-\mathbf{p}})} \leq \eta \|\mu_1 - \mu_2\|_{\mathcal{M}(1+\kappa V_{n-k})}.$$

### 3.2 Estimation of the eigenelements

Using the notations introduced in the previous section, we set now

$$\rho = \max \{ \eta, \mathfrak{a}^p \} = \max \left\{ 1 - (\mathfrak{b} - \mathfrak{b}'), \frac{2 + \kappa \mathfrak{R} \mathfrak{a}'}{2 + \kappa \mathfrak{R}}, \mathfrak{a}^p \right\} \in (0, 1), \quad (3.5)$$

with  $\mathfrak{b}' \in (0, \mathfrak{b})$  and  $\mathfrak{a}' \in (\mathfrak{a} + 2\mathfrak{c}/\mathfrak{R}, 1)$ . We first deal with the right eigenelement.

**Lemma 3.4.** *There exists  $h \in \mathcal{B}_+(V)$  and  $\lambda \in \mathbb{R}$  such that for all  $t \in [0, \infty)$ ,*

$$M_t h = e^{\lambda t} h.$$

Moreover,

$$\left( \frac{\psi}{V} \right)^q \psi \lesssim h \lesssim V, \quad \text{with } q = \frac{\log(cr)}{\log(\mathfrak{a})} > 0$$

and there exists  $C > 0$  such that for all integer  $k \geq 0$  and  $\mu \in \mathcal{M}_+(V)$ ,

$$\left| \mu(h) - \frac{\mu M_{k\tau} \psi}{\nu(M_{k\tau} \psi / \psi)} \right| \leq C \frac{\mu(V)^2}{\mu(\psi)} \rho^{\lfloor k/p \rfloor}. \quad (3.6)$$

Let us turn to the left eigenmeasure and provide a similar result.

**Lemma 3.5.** *There exists  $\gamma \in \mathcal{M}_+(V)$  such that  $\gamma(h) = 1$  and for all  $t \in [0, \infty)$ ,*

$$\gamma M_t = e^{\lambda t} \gamma.$$

Moreover, there exists  $C > 0$  such that for all integer  $k \geq 0$  and  $\mu \in \mathcal{M}_+(V)$ ,

$$\left\| \frac{\gamma(\cdot \psi)}{\gamma(\psi)} - \frac{\mu M_{k\tau}(\cdot \psi)}{\mu M_{k\tau}(\psi)} \right\|_{\mathcal{M}(1+\kappa V_0)} \leq C \left( \frac{\mu(V)}{\mu(\psi)} + \frac{\theta}{\beta - \alpha} \right) \rho^{\lfloor k/p \rfloor}. \quad (3.7)$$

Using Lemma 3.5,  $\psi \leq V$ , (A1) and (A2) yields  $e^{\lambda\tau} \gamma(V) = \gamma M_\tau V \leq (\alpha + \theta) \gamma(V)$  and  $e^{\lambda\tau} \gamma(\psi) = \gamma M_\tau \psi \geq \beta \gamma(\psi)$ . It gives the following estimate of the eigenvalue

$$\frac{\log(\beta)}{\tau} \leq \lambda \leq \frac{\log(\alpha + \theta)}{\tau}. \quad (3.8)$$

Theorem 2.1 follows from Lemmas 3.4 and 3.5. Indeed, existence and estimates on the eigenelements allow checking that  $(V, h)$  satisfies Assumption **A**. Then (3.7) with  $\psi$  replaced by  $h$  yields (2.2). Details are given in Section 4.5.

### 3.3 Uniform exponential convergence

We give here complementary results about the convergence to the profile given by the eigenelements. The first result provides a more explicit rate of convergence than in Theorem 2.1, to the price of loosing the  $V$ -uniformity. It requires the following additional assumption to capture the continuous time:

$$t \mapsto \sup_{x \in \mathcal{X}} \frac{M_t V(x)}{V(x)} \quad \text{and} \quad t \mapsto \sup_{x \in \mathcal{X}} \frac{\psi(x)}{M_t \psi(x)} \quad \text{are locally bounded on } [0, \infty). \quad (3.9)$$

**Theorem 3.6.** *Under Assumptions **A** and (3.9), there exists  $C > 0$  such that for all  $\mu \in \mathcal{M}_+(V)$  and  $t \geq 0$ ,*

$$\|\mu M_t - e^{\lambda t} \mu(h) \gamma\|_{\mathcal{M}(V)} \leq C \frac{\mu(V)}{\mu(\psi)} e^{-\sigma t} \min \{ \mu M_t \psi, \mu(V) e^{\lambda t} \},$$

where

$$\sigma = \frac{-\log \rho}{\mathbf{p}\tau} > 0.$$

Finally, the renormalisation of the semigroup by its mass  $M_t \mathbf{1}$  may also be relevant for applications. We mention the study of the convergence of the conditional probability to a quasi-stationary distribution and the study of the typical trait in a structured branching process, see respectively Section 5.1 and *e.g.* [8, 62, 61]. We write  $\mathcal{P}(V)$  for the set of probability measures which belong to  $\mathcal{M}(V)$  and we define the total variation norm for finite signed measures by  $\|\mu\|_{\text{TV}} = \|\mu\|_{\mathcal{M}(\mathbf{1})}$ . The following result is direct a corollary of Theorem 2.1.

**Corollary 3.7.** *Assume that conditions of Theorem 2.1-(i) hold and  $\inf_{\mathcal{X}} V > 0$ . Then there exist  $C, \omega > 0$  and  $\pi \in \mathcal{P}(V)$  such that for every  $\mu \in \mathcal{M}_+(V)$  and  $t \geq 0$ ,*

$$\left\| \frac{\mu M_t}{\mu M_t \mathbf{1}} - \pi \right\|_{\text{TV}} \leq C \frac{\mu(V)}{\mu(h)} e^{-\omega t}. \quad (3.10)$$

## 4 Proofs

### 4.1 Preliminary inequalities

For all  $t \geq 0$ , let us define the following operator

$$\widehat{M}_t : f \mapsto M_t(\mathbf{1}_{K^c} f),$$

and for convenience, we introduce the following constants

$$\Theta = \frac{\theta}{\beta - \alpha}, \quad R = \sup_K \frac{V}{\psi}, \quad \Xi = \alpha(R + \Theta) + \theta, \quad (4.1)$$

which are well-defined and finite under Assumption **A**. We first give some estimates which are directly deduced from Assumption **A**.

**Lemma 4.1.** *For all  $k \geq 0$ , we have*

$$i) \widehat{M}_\tau^k M_\tau V \leq \alpha^k M_\tau V,$$

ii) *for all  $\mu \in \mathcal{M}_+(V)$ ,*

$$\frac{\mu M_{k\tau} V}{\mu M_{k\tau} \psi} \leq \mathbf{a}^k \frac{\mu(V)}{\mu(\psi)} + \Theta,$$

iii) *for all  $x \in K$  and  $n \geq k$ ,  $M_{n\tau} \psi(x) \leq \Xi^k M_{(n-k)\tau} \psi(x)$ ,*

iv) *for all  $x \in K$ , and  $f \in \mathcal{B}_+(V/\psi)$ ,*

$$M_{(k+1)\tau}(f\psi)(x) \geq c_{k+1} \nu(f) M_{(k+1)\tau} \psi(x), \quad \text{with } c_{k+1} = c^{k+1} (\beta/\Xi)^k. \quad (4.2)$$

**Remark 4.2.** We observe that  $(c_k)_{k \geq 1}$  is a decreasing geometric sequence. Indeed since  $\psi \leq V$ , (A1) and (A2) ensure that on  $K$ ,  $\beta\psi \leq M_\tau \psi \leq M_\tau V \leq (\alpha R + \theta)\psi$ , so that  $\beta < \Xi$ . Adding that  $c < 1$  ensures that  $(c_k)_{k \geq 1}$  decreases geometrically.

Points *i)* and *ii)* of Lemma 4.1 are sharp inequalities, while *iv)* extends Assumption (A3) for any time.

*Proof.* Using (A1) we readily have  $\mathbf{1}_{K^c} M_\tau V \leq \alpha V$  and *i)* follows by induction. Composing respectively (A1) and (A2) with  $M_{k\tau}$  yields

$$M_{(k+1)\tau} V \leq \alpha M_{k\tau} V + \theta M_{k\tau} \psi; \quad M_{(k+1)\tau} \psi \geq \beta M_{k\tau} \psi.$$

Combining these inequalities gives

$$\frac{M_{(k+1)\tau} V}{M_{(k+1)\tau} \psi} \leq \mathbf{a} \frac{M_{k\tau} V}{M_{k\tau} \psi} + \frac{\theta}{\beta}$$

and *ii)* follows by induction recalling that  $\mathbf{a} < 1$ .

By definition of  $R$ , we immediately deduce from *ii)* that for any  $x \in K$ ,

$$\frac{M_{k\tau} V(x)}{M_{k\tau} \psi(x)} \leq R + \Theta.$$

Combining this inequality with

$$M_{n\tau} \psi \leq M_{(n-1)\tau} M_\tau V \leq M_{(n-1)\tau} (\alpha V + \theta \psi),$$

coming from (A1) and  $\psi \leq V$ , yields for  $x \in K$ ,  $M_{n\tau} \psi(x) \leq \Xi M_{(n-1)\tau} \psi(x)$ . The proof of *iii)* is completed by induction.



Finally, let  $x \in K$ . We have

$$\frac{M_{(n+1)\tau}(f\psi)(x)}{M_{(n+1)\tau}\psi(x)} = \frac{M_\tau(M_{n\tau}(f\psi))(x)}{M_{(n+1)\tau}\psi(x)} \geq c\nu\left(\frac{M_{n\tau}(f\psi)}{\psi}\right) \frac{M_\tau\psi(x)}{M_{(n+1)\tau}\psi(x)},$$

using (A3) with the function  $M_{n\tau}(f\psi)/\psi$ . Besides, since  $\nu$  is supported by  $K$ , (A3) and (A2) yield

$$\begin{aligned} \nu\left(\frac{M_{n\tau}(f\psi)}{\psi}\right) &= \nu\left(\frac{M_\tau(M_{(n-1)\tau}(f\psi))}{\psi}\right) \\ &\geq c\nu\left(\nu\left(\frac{M_{(n-1)\tau}(f\psi)}{\psi}\right) \frac{M_\tau\psi}{\psi}\right) \geq c\beta\nu\left(\frac{M_{(n-1)\tau}(f\psi)}{\psi}\right). \end{aligned}$$

Iterating the last inequality and plugging it in the previous one, we obtain

$$\frac{M_{(n+1)\tau}(f\psi)(x)}{M_{(n+1)\tau}\psi(x)} \geq c^{n+1}\beta^n \frac{M_\tau\psi(x)}{M_{(n+1)\tau}\psi(x)} \nu(f).$$

Moreover, for  $x \in K$ , we have  $M_{(n+1)\tau}\psi(x) \leq \Xi^n M_\tau\psi(x)$  using *iii*). We obtain

$$\frac{M_{(n+1)\tau}(f\psi)(x)}{M_{(n+1)\tau}\psi(x)} \geq c^{n+1} \frac{\beta^n}{\Xi^n} \nu(f),$$

for all  $x \in K$ , which completes the proof.  $\square$

## 4.2 Contraction property: proofs of Section 3.1

First, we prove that  $(V_k)_{k \geq 0}$  is a family of Lyapunov functions for the sequence of operators  $(P_{k\tau, (k+1)\tau}^{(n\tau)})_{0 \leq k \leq n-1}$ .

**Lemma 4.3.** *For all  $k \geq 0$  and  $n \geq m \geq k$ , we have*

$$P_{k\tau, m\tau}^{(n\tau)} V_{n-m} \leq \mathbf{a}^{m-k} V_{n-k} + \frac{\theta}{c\beta} \sum_{j=k}^{m-1} \mathbf{a}^{m-j} P_{k\tau, j\tau}^{(n\tau)} (\mathbf{1}_K).$$

*Proof.* By definition of  $V_k$  in (3.2), we have, for  $0 \leq k \leq n$ ,

$$P_{(k-1)\tau, k\tau}^{(n\tau)} V_{n-k} = \frac{M_\tau(V_{n-k} M_{(n-k)\tau}\psi)}{M_{(n-k+1)\tau}\psi} = \nu\left(\frac{M_{(n-k)\tau}\psi}{\psi}\right) \frac{M_\tau V}{M_{(n-k+1)\tau}\psi}$$

Using (A1) and (A2), we have  $M_\tau V \leq \alpha V + \theta\psi\mathbf{1}_K$  and  $M_{(n-k)\tau}\psi \leq M_{(n-k+1)\tau}\psi/\beta$ . We obtain from the definitions of  $\mathbf{a}$  and  $V_{n-k+1}$  that

$$P_{(k-1)\tau, k\tau}^{(n\tau)} V_{n-k} \leq \mathbf{a} V_{n-k+1} + \nu\left(\frac{M_{(n-k)\tau}\psi}{\psi}\right) \frac{\theta\psi\mathbf{1}_K}{M_{(n-k+1)\tau}\psi}.$$

Besides, combining (A2) and (A3) with  $f = M_{(n-k)\tau}\psi/\psi$ , we get

$$\nu \left( \frac{M_{(n-k)\tau}\psi}{\psi} \right) \frac{\psi \mathbf{1}_K}{M_{(n-k+1)\tau}\psi} \leq \frac{\mathbf{1}_K}{c\beta}.$$

The last two inequalities yield

$$P_{(k-1)\tau, k\tau}^{(n\tau)} V_{n-k} \leq \mathbf{a} V_{n-k+1} + \frac{\theta \mathbf{1}_K}{c\beta}.$$

The conclusion follows from  $P_{k\tau, m\tau}^{(n\tau)} V_{n-m} = P_{k\tau, (k+1)\tau}^{(n\tau)} \cdots P_{(m-1)\tau, m\tau}^{(n\tau)} V_{n-m}$ .  $\square$

*Proof of Lemma 3.1.* Using that  $P_{k\tau, (j-1)\tau}^{(n\tau)}(\mathbf{1}_K) \leq 1$  and  $\mathbf{a} < 1$ , it is a direct consequence of Lemma 4.3.  $\square$

Using (A3) and (A4) and following [21, 8], we prove a small set condition (6.2) on the set  $K$  for the embedded semigroup  $P^{(t)}$ . However, Theorem 6.1 requires that (6.2) is satisfied on a sublevel set of  $V_k$ . Although there exists  $R > 0$  such that  $K \subset \{V_k \leq R\}$ , nothing guarantees the other inclusion. This situation is reminiscent of [48, Assumption 3] and we adapt here their arguments. For that purpose, we need a lower bound for the Lyapunov functions  $(V_k)_{k \geq 0}$ , which is stated in the next lemma.

**Lemma 4.4.** *For every  $n \geq 0$ , we have*

$$d_1 M_{(n+1)\tau}\psi \leq \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) M_\tau V \quad \text{and} \quad V_n \geq d_2,$$

with  $d_1 = (1 - \mathbf{a})d$ ,  $d_2 = (\beta - \alpha)d/(\alpha + \theta)$ .

*Proof.* First, using (A4),

$$d M_{(n+1)\tau}\psi = d M_\tau (\mathbf{1}_K M_{n\tau}\psi + \mathbf{1}_{K^c} M_{n\tau}\psi) \leq \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) M_\tau \psi + d \widehat{M}_\tau M_{n\tau}\psi.$$

Then, by iteration, using (A2) and  $\psi \leq V$ ,

$$d M_{(n+1)\tau}\psi \leq \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) \sum_{j=0}^n \beta^{-j} \widehat{M}_\tau^j M_\tau \psi \leq \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) \sum_{j=0}^n \beta^{-j} \widehat{M}_\tau^j M_\tau V.$$

Hence by Lemma 4.1 i)

$$d M_{(n+1)\tau}\psi \leq \frac{1}{1 - \mathbf{a}} \nu \left( \frac{M_{n\tau}\psi}{\psi} \right) M_\tau V$$

and the first identity is proved. From the definition of  $V_n$  we deduce

$$V_n \geq d_1 \frac{M_{(n+1)\tau}\psi}{M_{n\tau}\psi} \frac{V}{M_\tau V} \geq d_1 \frac{\beta}{\alpha + \theta} = d_2$$

by using successively (A2), (A1), and  $\psi \leq V$ .  $\square$

We now prove the small set condition (6.2) for the embedded semigroup.

*Proof of Lemma 3.2.* First, we introduce the measure  $\nu_i$  defined by

$$\nu_i(f) = \nu \left( f \frac{M_{i\tau}\psi}{\psi} \right)$$

for all integer  $i \geq 0$ . For any  $x \in K$ ,  $j \leq k \leq n$ , we have using Lemma 4.1 *iv*) with the function  $fM_{(n-k)\tau}\psi/\psi$ ,

$$P_{(j-1)\tau, k\tau}^{(n\tau)} f(x) = \frac{M_{(k-j+1)\tau} \left( fM_{(n-k)\tau}\psi \right) (x)}{M_{(n-j+1)\tau}\psi(x)} \geq c_{k-j+1} \nu_{n-k}(f) \frac{M_{(k-j+1)\tau}\psi(x)}{M_{(n-j+1)\tau}\psi(x)}.$$

for any nonnegative measurable function  $f$ . Then, Lemma 4.4 and (A2) yield

$$P_{(j-1)\tau, k\tau}^{(n\tau)} f(x) \geq d_1 c_{k-j+1} \frac{\nu_{n-k}(f)}{\nu_{n-j}(\mathbf{1})} \frac{M_{(k-j+1)\tau}\psi(x)}{M_\tau V(x)} \geq d_1 c_{k-j+1} \beta^{k-j} \frac{\nu_{n-k}(f)}{\nu_{n-j}(\mathbf{1})} \frac{M_\tau\psi(x)}{M_\tau V(x)}.$$

Recalling from (A1) and (A2) that for  $x \in K$ ,  $M_\tau\psi(x)/M_\tau V(x) \geq \beta/(\alpha R + \theta)$ , and from Lemma 4.1 *iii*) and  $\nu(K) = 1$  that  $\nu_{n-k}(\mathbf{1})/\nu_{n-j}(\mathbf{1}) \geq \Xi^{-(k-j)}$ , we get

$$P_{(j-1)\tau, k\tau}^{(n\tau)} f(x) \geq \alpha_{k-j} \frac{\nu_{n-k}(f)}{\nu_{n-k}(\mathbf{1})}, \quad (4.3)$$

for  $x \in K$ , where  $\alpha_i = d_1 c^{i+1} \beta r^i / (\alpha R + \theta)$  and  $r = (\beta/\Xi)^2$ . The previous bound holds only on  $K$ . We prove now that the semigroup charges  $K$  at an intermediate time and derive the expected lower bound. More precisely, setting

$$\omega_i = \frac{\mathbf{a}^i}{\alpha_i} \quad \text{and} \quad S_\ell = \sum_{j=1}^{\ell} \omega_{\ell-j} = \frac{\alpha R + \theta}{dc(\beta - \alpha)} \cdot \sum_{j=1}^{\ell} \left( \frac{\mathbf{a}}{cr} \right)^j,$$

we obtain for  $k \leq n - 1$  and  $1 \leq \ell \leq n - k$ ,

$$\begin{aligned} P_{k\tau, (k+\ell)\tau}^{(n\tau)} f &= \frac{1}{S_\ell} \sum_{j=k+1}^{k+\ell} \omega_{k+\ell-j} P_{k\tau, (j-1)\tau}^{(n\tau)} P_{(j-1)\tau, (k+\ell)\tau}^{(n\tau)}(f) \\ &\geq \frac{1}{S_\ell} \sum_{j=k+1}^{k+\ell} \omega_{k+\ell-j} P_{k\tau, (j-1)\tau}^{(n\tau)} (\mathbf{1}_K P_{(j-1)\tau, (k+\ell)\tau}^{(n\tau)} f) \geq B_{k,n}^{(\ell)} \frac{\nu_{n-k-\ell}(f)}{\nu_{n-k-\ell}(\mathbf{1})}, \end{aligned}$$

where the last inequality comes from (4.3) and

$$B_{k,n}^{(\ell)} = \frac{1}{S_\ell} \sum_{j=k+1}^{k+\ell} \mathbf{a}^{k+\ell-j} P_{k\tau, (j-1)\tau}^{(n\tau)} \mathbf{1}_K.$$

To conclude, we need to find a positive lower bound for  $B_{k,n}^{(\ell)}$  which does not depend on  $k$  or  $n$ . For that purpose, we first observe that the second bound of Lemma 4.4 ensures that  $P_{k\tau, (k+\ell)\tau}^{(n\tau)} V_{n-k-\ell} \geq d_2$ . Using now Lemma 4.3 yields

$$\sum_{j=k+1}^{k+\ell} \mathbf{a}^{k+\ell-j} P_{k\tau, (j-1)\tau}^{(n\tau)} \mathbf{1}_K \geq c\beta \frac{d_2 - \mathbf{a}^\ell V_{n-k}}{\theta},$$

for  $n \geq k + \ell$ . For  $x \in \{V_{n-k} \leq \mathfrak{R}\}$  and  $\ell = \mathfrak{p}$  defined in (3.4), we get

$$B_{k,n}^{(\mathfrak{p})}(x) \geq c\beta \frac{d_2}{2\theta S_{\mathfrak{p}}} = \frac{c^2 \beta^3 d_1^2}{2\theta(\alpha + \theta)(\alpha R + \theta)} \frac{1}{\sum_{j=1}^{\mathfrak{p}} (\mathbf{a}/c\mathfrak{r})^j},$$

which ends the proof.  $\square$

*Proof of Proposition 3.3.* Let  $n$  and  $k$  be two integers such that  $0 \leq k \leq n - \mathfrak{p}$  and consider  $\mathfrak{R} > 2\mathfrak{c}/(1 - \mathbf{a})$ . According to Lemmas 3.1 and 3.2, the conservative operator  $P_{k\tau, (k+\mathfrak{p})\tau}^{(n\tau)}$  satisfies condition (6.1) with the functions  $V_{n-k-\mathfrak{p}}$  and  $V_{n-k}$  and condition (6.2) with the probability measure  $\nu_{k,n}$ . Applying Theorem 6.1 then yields the contraction result.  $\square$

### 4.3 Eigenelements: proofs of Section 3.2

Let us consider, for every  $\mu \in \mathcal{M}_+(V)$ , the family of operators  $(Q_t^\mu)_{t \geq 0}$  defined for  $f \in \mathcal{B}(V_0)$  by

$$Q_t^\mu f = \frac{\mu M_t(\psi f)}{\mu M_t(\psi)}.$$

Fixing the measure  $\mu$ , the operator  $f \mapsto Q_t^\mu f$  is linear. Observe that  $Q_t^{\delta_x} = \delta_x P_{0,t}^{(t)}$  so that Proposition 3.3 implies contraction inequalities for  $\delta_x \mapsto Q_{n\mathfrak{p}\tau}^{\delta_x}$ . Notice that  $\mu \mapsto Q_{n\mathfrak{p}\tau}^\mu$  is non-linear and forthcoming Lemma 4.6 extends the contraction to a more general space of measures. Besides, for any positive measure  $\mu$ , we set

$$[\mu] = \frac{\mu(V)}{\mu(\psi)}.$$

We prove now the existence of the eigenvector and eigenmeasure, respectively stated in Lemma 3.4 and Lemma 3.5. Let us first provide a useful upper bound for  $V_k$ . For that purpose, we also set

$$p = \left\lceil \frac{\log(2(1 + \theta/\alpha)(\Theta + R))}{\log(1/\mathbf{a})} \right\rceil + 1, \quad C_1 = \frac{2\Xi^{p+1}}{cc_{p-1}\beta^{p+1}}. \quad (4.4)$$

where  $(c_k)_{k \geq 0}$  is defined in (4.2).

**Lemma 4.5.** *For all positive measure  $\mu$  such that*

$$[\mu] \leq \Theta + R, \quad (4.5)$$

*we have for all  $k \geq p$ ,*

$$\nu \left( \frac{M_{k\tau}\psi}{\psi} \right) \leq C_1 \frac{\mu M_{k\tau}\psi}{\mu(\psi)}. \quad (4.6)$$

The idea is the following: condition (4.5) ensures the existence of a time  $p$  at which the semigroup charges  $K$ . Then, (A3) yields (4.6). It will be needed in this form in the sequel, but could be extended to more general right-hand side in (4.5).

*Proof.* Recalling that  $\widehat{M}_\tau = M_\tau(\mathbf{1}_{K^c} \cdot)$  and using that for all  $g \in \mathcal{B}(V)$ ,  $M_{(k+1)\tau}g = M_\tau(\mathbf{1}_K M_{k\tau}g) + \widehat{M}_\tau(M_{k\tau}g)$ , we obtain by induction

$$M_{k\tau}g = \widehat{M}_\tau^k g + \sum_{j=1}^k \widehat{M}_\tau^{k-j} M_\tau(\mathbf{1}_K M_{(j-1)\tau}g).$$

Let  $g = \psi \mathbf{1}_K$ . Using Lemma 4.1 *iv*) with  $f = \mathbf{1}_K$  and that  $\nu(K) = 1$ , we have

$$\begin{aligned} M_{k\tau}(\mathbf{1}_K\psi) &\geq \sum_{j=1}^k \widehat{M}_\tau^{k-j} M_\tau(\mathbf{1}_K M_{(j-1)\tau}(\mathbf{1}_K\psi)) \\ &\geq \sum_{j=1}^k c_{j-1} \widehat{M}_\tau^{k-j} M_\tau(\mathbf{1}_K M_{(j-1)\tau}\psi) \\ &\geq c_{k-1} \sum_{j=1}^k \left( \widehat{M}_\tau^{k-j} M_{j\tau}\psi - \widehat{M}_\tau^{k-j} M_\tau(\mathbf{1}_{K^c} M_{(j-1)\tau}\psi) \right) \\ &= c_{k-1} \left( M_{k\tau}\psi - \widehat{M}_\tau^k \psi \right), \end{aligned}$$

with the convention that  $c_0 = 1$ . Then, using (A2) and the fact that  $\widehat{M}_\tau^k \psi \leq \widehat{M}_\tau^{k-1} M_\tau V$  together with Lemma 4.1 *i*) yields  $M_{k\tau}(\mathbf{1}_K\psi) \geq c_{k-1} (\beta^k \psi - \alpha^{k-1} M_\tau V)$ . Next, (A1) and the fact that  $V \geq \psi$  yield

$$M_{k\tau}(\mathbf{1}_K\psi) \geq c_{k-1} \beta^k (\psi - \mathbf{a}^k (1 + \theta/\alpha) V). \quad (4.7)$$

Using the definition (4.4) of  $p$  and the fact that  $\mathbf{a} \leq 1$  ensure that  $\mathbf{a}^{p2} (1 + \theta/\alpha) (\Theta + R) \leq 1$ , and (4.5) yields  $\mathbf{a}^p (1 + \theta/\alpha) \mu(V) \leq \mu(\psi)/2$ . Then, (4.7) becomes

$$\mu M_{p\tau}(\mathbf{1}_K\psi) \geq c_{p-1} \beta^p \mu(\psi)/2. \quad (4.8)$$

Using  $\mu M_{n\tau}\psi \geq \mu M_{p\tau}(\mathbf{1}_K M_{(n-p)\tau}\psi) = \mu M_{p\tau}(\mathbf{1}_K M_\tau((M_{(n-p-1)\tau}\psi/\psi)\psi))$  for  $n \geq p$  and successively (A3) with  $f = M_{(n-p-1)\tau}\psi/\psi$ , (A2) and (4.8), we get

$$\mu M_{n\tau}\psi \geq c c_{p-1} \beta^{p+1} \frac{\mu(\psi)}{2} \nu \left( \frac{M_{(n-p-1)\tau}\psi}{\psi} \right).$$

Finally, combining this estimate with Lemma 4.1 *iii*) ensures that

$$\nu \left( \frac{M_{n\tau}\psi}{\psi} \right) \leq \Xi^{p+1} \nu \left( \frac{M_{(n-p-1)\tau}\psi}{\psi} \right) \leq C_1 \frac{\mu M_{n\tau}\psi}{\mu(\psi)},$$

which ends the proof.  $\square$

We generalize now Proposition 3.3 to  $(Q_{n\mathfrak{p}\tau}^\mu)_{n \geq 0}$ . Recall that  $\mathfrak{p}$  is defined in Lemma 3.2,  $\kappa$  and  $\mathfrak{v}$  are defined in Proposition 3.3 and  $\rho$  is defined in (3.5).

**Lemma 4.6.** *For all measures  $\mu_1, \mu_2 \in \mathcal{M}_+(V/\psi)$  and all  $n \geq 0$  we have*

$$\|Q_{n\mathfrak{p}\tau}^{\mu_1} - Q_{n\mathfrak{p}\tau}^{\mu_2}\|_{\mathcal{M}(1+\kappa V_0)} \leq C_2 \rho^n ([\mu_1] + [\mu_2]), \quad (4.9)$$

where

$$C_2 = \max \left\{ 2\mathfrak{a}^{-\mathfrak{p}} + \kappa C_1 (1 + 2\Theta \mathfrak{a}^{-\mathfrak{p}}), 2(1 + \kappa \Theta) \mathfrak{a}^{-(p+\mathfrak{p})} + \kappa \right\}$$

with  $C_1$  and  $p$  defined in (4.4).

*Proof.* Fix  $\mu_1, \mu_2 \in \mathcal{M}_+(V/\psi)$ ,  $f \in \mathcal{B}(V/\psi)$  with  $\|f\|_{\mathcal{B}(1+\kappa V_0)} \leq 1$  and an integer  $n \geq 0$ . Set for convenience

$$m = \left\lfloor \frac{\log([\mu_1] + [\mu_2])}{\mathfrak{p} \log(1/\mathfrak{a})} \right\rfloor + 1, \quad \mathfrak{n} = \mathfrak{p}\tau n, \quad \mathfrak{m} = \mathfrak{p}\tau m. \quad (4.10)$$

By definition of the embedded semigroup in (3.1), we have

$$\begin{aligned} & \mu_1 M_{\mathfrak{n}}(f\psi) \mu_2 M_{\mathfrak{n}}\psi - \mu_2 M_{\mathfrak{n}}(f\psi) \mu_1 M_{\mathfrak{n}}\psi \\ &= \int_{\mathcal{X}^2} M_{\mathfrak{n}}\psi(x) M_{\mathfrak{n}}\psi(y) \left( \frac{M_{\mathfrak{n}}(f\psi)(x)}{M_{\mathfrak{n}}\psi(x)} - \frac{M_{\mathfrak{n}}(f\psi)(y)}{M_{\mathfrak{n}}\psi(y)} \right) \mu_1(dx) \mu_2(dy) \\ &\leq \int_{\mathcal{X}^2} M_{\mathfrak{n}}\psi(x) M_{\mathfrak{n}}\psi(y) \left\| \delta_x P_{0,\mathfrak{n}}^{(\mathfrak{n})} - \delta_y P_{0,\mathfrak{n}}^{(\mathfrak{n})} \right\|_{\mathcal{M}(1+\kappa V_0)} \mu_1(dx) \mu_2(dy). \end{aligned}$$

Using Proposition 3.3, we get for  $n \geq m$ ,

$$\begin{aligned} & \mu_1 M_{\mathfrak{n}}(f\psi) \mu_2 M_{\mathfrak{n}}\psi - \mu_2 M_{\mathfrak{n}}(f\psi) \mu_1 M_{\mathfrak{n}}\psi \\ &\leq \mathfrak{v}^{n-m} \int_{\mathcal{X}^2} M_{\mathfrak{n}}\psi(x) M_{\mathfrak{n}}\psi(y) \left\| \delta_x P_{0,\mathfrak{m}}^{(\mathfrak{n})} - \delta_y P_{0,\mathfrak{m}}^{(\mathfrak{n})} \right\|_{\mathcal{M}(1+\kappa V_{(n-m)\mathfrak{p}})} \mu_1(dx) \mu_2(dy). \end{aligned} \quad (4.11)$$

Using the definition of the norm on  $\mathcal{M}(1 + \kappa V_{(n-m)\mathfrak{p}})$  and the definition of  $V_{(n-m)\mathfrak{p}}$  in (3.2), we obtain

$$\begin{aligned} \left\| \delta_x P_{0,\mathfrak{m}}^{(\mathfrak{n})} - \delta_y P_{0,\mathfrak{m}}^{(\mathfrak{n})} \right\|_{\mathcal{M}(1+\kappa V_{(n-m)\mathfrak{p}})} &\leq \int_{\mathcal{X}} (1 + \kappa V_{(n-m)\mathfrak{p}}(z)) \left| \delta_x P_{0,\mathfrak{m}}^{(\mathfrak{n})} - \delta_y P_{0,\mathfrak{m}}^{(\mathfrak{n})} \right| (dz) \\ &\leq 2 + \kappa \nu \left( \frac{M_{n-m}\psi}{\psi} \right) \left( \frac{M_{\mathfrak{m}}V(x)}{M_{\mathfrak{n}}\psi(x)} + \frac{M_{\mathfrak{m}}V(y)}{M_{\mathfrak{n}}\psi(y)} \right). \end{aligned}$$

Combining this inequality with (4.11) and  $\rho \geq \mathfrak{h}$ , we get

$$\begin{aligned} Q_n^{\mu_1} f - Q_n^{\mu_2} f &= \frac{\mu_1 M_n(f\psi) \mu_2 M_n \psi - \mu_2 M_n(f\psi) \mu_1 M_n \psi}{\mu_1 M_n \psi \cdot \mu_2 M_n \psi} \\ &\leq \rho^n \left( 2\rho^{-m} + \kappa \rho^{-m} \nu \left( \frac{M_{n-m}\psi}{\psi} \right) \left( \frac{\mu_1 M_m V}{\mu_1 M_n \psi} + \frac{\mu_2 M_m V}{\mu_2 M_n \psi} \right) \right). \end{aligned} \quad (4.12)$$

We now bound each term of the right-hand side. First, using that  $\mathfrak{a}^p \leq \rho$  and (4.10),

$$\mathfrak{a}^p \leq \rho^m ([\mu_1] + [\mu_2]). \quad (4.13)$$

Second, Lemma 4.1 *ii*) ensures that for  $\mu \in \{\mu_1, \mu_2\}$ ,

$$[\mu M_m] \leq \mathfrak{a}^{mp} [\mu] + \Theta. \quad (4.14)$$

Besides (4.10) also guarantees that for  $\mu \in \{\mu_1, \mu_2\}$ ,  $\mathfrak{a}^{mp} [\mu] \leq 1$ . It means that the positive measure  $\mu M_m$  satisfies inequality (4.5), since  $R \geq 1$ . Then, Lemma 4.5 applied to  $\mu M_m$  with  $k = (n - m)\mathfrak{p}$  yields for all  $n \geq m + p/\mathfrak{p}$ ,

$$\nu \left( \frac{M_{n-m}\psi}{\psi} \right) \frac{\mu M_m V}{\mu M_n \psi} \leq C_1 \frac{\mu M_m M_{n-m}\psi}{\mu M_m \psi} \frac{\mu M_m V}{\mu M_n \psi} \leq C_1 [\mu M_m].$$

Finally, using again (4.14) and (4.13), we get

$$\nu \left( \frac{M_{n-m}\psi}{\psi} \right) \left( \frac{\mu_1 M_m V}{\mu_1 M_n \psi} + \frac{\mu_2 M_m V}{\mu_2 M_n \psi} \right) \leq C_1 (1 + 2\Theta \mathfrak{a}^{-p}) \rho^m ([\mu_1] + [\mu_2]).$$

Plugging the last inequality in (4.12) ensures that for all  $n \geq m + p/\mathfrak{p}$ ,

$$Q_n^{\mu_1} f - Q_n^{\mu_2} f \leq (2\mathfrak{a}^{-p} + \kappa C_1 (1 + 2\Theta \mathfrak{a}^{-p})) ([\mu_1] + [\mu_2]) \rho^n.$$

To conclude, it remains to show that (4.9) also holds for  $n \leq m + p/\mathfrak{p}$ . We have

$$\|Q_n^{\mu_1} - Q_n^{\mu_2}\|_{\mathcal{M}(1+\kappa V_0)} \leq \|Q_n^{\mu_1}\|_{\mathcal{M}(1+\kappa V_0)} + \|Q_n^{\mu_2}\|_{\mathcal{M}(1+\kappa V_0)} \leq 2 + \kappa [\mu_1 M_n] + \kappa [\mu_2 M_n].$$

Using again (4.14),  $[\mu M_n] \leq \mathfrak{a}^{np} [\mu] + \Theta \leq \rho^n [\mu] + \Theta$  for  $\mu \in \{\mu_1, \mu_2\}$ , so that

$$\|Q_n^{\mu_1} - Q_n^{\mu_2}\|_{\mathcal{M}(1+\kappa V_0)} \leq 2(1 + \kappa \Theta) + \kappa \rho^n ([\mu_1] + [\mu_2]).$$

Finally,  $\rho \geq \mathfrak{a}^p$  and  $n \leq m + p/\mathfrak{p}$  and (4.10) yield  $1 \leq \rho^n \mathfrak{a}^{-(p+mp)} = \rho^n \mathfrak{a}^{-(p+p)} \mathfrak{a}^{-(m-1)\mathfrak{p}} \leq \rho^n \mathfrak{a}^{-(p+p)} ([\mu_1] + [\mu_2])$ , and we get

$$\|Q_n^{\mu_1} - Q_n^{\mu_2}\|_{\mathcal{M}(1+\kappa V_0)} \leq \rho^n (2(1 + \kappa \Theta) \mathfrak{a}^{-(p+p)} + \kappa) ([\mu_1] + [\mu_2]),$$

for all  $n \leq m + p/\mathfrak{p}$ , which ends the proof.  $\square$

We have now all the ingredients to prove the existence of the eigenelements and the associated estimates. We start with the right eigenfunction.

*Proof of Lemma 3.4.* We define

$$\eta(\cdot) = \nu(\cdot/\psi)$$

and, for  $k \geq 0$ ,

$$h_k = \frac{M_{k\tau}\psi}{\nu(M_{k\tau}\psi/\psi)} = \frac{M_{k\tau}\psi}{\eta M_{k\tau}\psi}.$$

The proof is divided into four steps. We begin by giving preliminary estimates on  $M_{k\tau}\psi$ . Next, we show that  $h_k$  converges in  $\mathcal{B}(V^2/\psi)$  as  $k \rightarrow \infty$  and that its limit  $h$  satisfies

$$h \leq d_2^{-1}V, \quad (4.15)$$

where  $d_2$  is defined in Lemma 4.4. Then, we show that  $h$  is an eigenvector and we give a lower bound for  $h$ .

Using the first part of Lemma 4.4 and (A1), we obtain for any  $k \geq 0$ ,

$$M_{(k+1)\tau}\psi(x) \leq d_1^{-1}(\alpha + \theta)(\eta M_{k\tau}\psi)V(x). \quad (4.16)$$

Since  $\|M_{k\tau}\psi/\psi\|_{\mathcal{B}(1+\kappa V_0)} = \sup_{\mathcal{X}} M_{k\tau}\psi/(\psi + \kappa V)$ , we get for any  $k \geq 0$ ,

$$\left\| \frac{M_{(k+1)\tau}\psi}{\psi} \right\|_{\mathcal{B}(1+\kappa V_0)} \leq d_1^{-1}(\alpha + \theta)(\eta M_{k\tau}\psi) \sup_{\mathcal{X}} \frac{V}{\psi + \kappa V} \leq \frac{(\alpha + \theta)}{d_1 \kappa} \eta M_{k\tau}\psi. \quad (4.17)$$

Besides, from Lemma 4.1 *ii*) and the fact that  $V \leq R\psi$  on  $K$ ,  $[\eta M_{k\tau}] \leq \mathfrak{a}^k \eta(V) + \Theta \leq R + \Theta$ , so  $\mu = \eta M_{k\tau}$  verifies (4.5). Lemma 4.5 applied to  $\mu$  gives for all  $n \geq p$ ,

$$(\eta M_{k\tau})(M_{n\tau}\psi) \geq C_1^{-1}(\eta M_{n\tau}\psi)(\eta M_{k\tau}\psi). \quad (4.18)$$

We can now proceed to the second step: the convergence of  $(h_k)_{k \geq 0}$ . Let  $\mu \in \mathcal{M}_+(V)$ . We use that

$$\mu(h_{k+n}) = \frac{\mu M_{(k+n)\tau}\psi / \mu M_{k\tau}\psi}{\eta M_{(k+n)\tau}\psi / \eta M_{k\tau}\psi} \mu(h_k)$$

to obtain that

$$\begin{aligned} |\mu(h_{k+n}) - \mu(h_k)| &\leq \left| \frac{\mu M_{(k+n)\tau}\psi}{\mu M_{k\tau}\psi} - \frac{\eta M_{(k+n)\tau}\psi}{\eta M_{k\tau}\psi} \right| \frac{1}{\eta M_{(k+n)\tau}\psi / \eta M_{k\tau}\psi} \mu(h_k) \\ &= \left| Q_{k\tau}^\mu \left( \frac{M_{n\tau}\psi}{\psi} \right) - Q_{k\tau}^\eta \left( \frac{M_{n\tau}\psi}{\psi} \right) \right| \frac{\mu M_{k\tau}\psi}{\eta M_{(k+n)\tau}\psi}. \end{aligned}$$

Then, Lemma 4.6 yields

$$|\mu(h_{k+n}) - \mu(h_k)| \leq C_2 \rho^{\lfloor k/p \rfloor} ([\mu M_{m\tau}] + [\eta M_{m\tau}]) \left\| \frac{M_{n\tau}\psi}{\psi} \right\|_{\mathcal{B}(1+\kappa V_0)} \frac{\mu M_{k\tau}\psi}{\eta M_{(k+n)\tau}\psi}, \quad (4.19)$$



where  $m = k - \lfloor k/\mathfrak{p} \rfloor \mathfrak{p}$ . First, using (A1) and (A2) we have since  $m \leq \mathfrak{p}$

$$[\mu M_{m\tau}] \leq \max \left( 1, \frac{(\alpha + \theta)^\mathfrak{p}}{\beta^\mathfrak{p}} \right) [\mu]. \quad (4.20)$$

Next, combining (4.19) with (4.16), (4.17), (4.18), (4.20), and using (A2) yields

$$|\mu(h_{k+n}) - \mu(h_k)| \leq C_1 C_2' \frac{(\alpha + \theta)^2}{d_1^2 \beta^2 \kappa} \rho^{\lfloor k/\mathfrak{p} \rfloor} \left( [\mu] + \nu \left( \frac{V}{\psi} \right) \right) \mu(V),$$

for all  $n \geq p$ , where

$$C_2' = C_2 \max \left( 1, \frac{(\alpha + \theta)^\mathfrak{p}}{\beta^\mathfrak{p}} \right). \quad (4.21)$$

Recalling that  $\psi \leq V$ ,  $V \leq R\psi$  on  $K$  and  $\nu(K^c) = 0$ , we get for all  $n \geq p$

$$|\mu(h_{k+n}) - \mu(h_k)| \leq C_3 \rho^{\lfloor k/\mathfrak{p} \rfloor} \mu(V) [\mu], \quad (4.22)$$

with  $C_3 = C_1 C_2' \frac{(\alpha + \theta)^2}{d_1^2 \beta^2 \kappa} (1 + R)$ . Taking  $\mu = \delta_x$  we deduce for all  $n \geq p$

$$\|h_{k+n} - h_k\|_{\mathcal{B}(V^2/\psi)} \leq C_3 \rho^{\lfloor k/\mathfrak{p} \rfloor}.$$

Cauchy criterion ensures that  $h_k$  converges as  $k \rightarrow \infty$  in  $\mathcal{B}(V^2/\psi)$  to a limit denoted by  $h$ . Moreover from Lemma 4.4,  $d_2 \leq V_m = V/h_{m\tau}$  which yields (4.15) by letting  $m \rightarrow \infty$ . Letting  $n \rightarrow \infty$  in (4.22) gives (3.6).

We move on to the third step and check that  $h$  is an eigenfunction. For all  $x \in \mathcal{X}$ ,

$$\eta M_\tau \left( \frac{M_{n\tau} \psi}{\eta M_{n\tau} \psi} \right) \frac{M_{(n+1)\tau} \psi(x)}{\eta M_{(n+1)\tau} \psi} = \delta_x M_\tau \left( \frac{M_{n\tau} \psi}{\eta M_{n\tau} \psi} \right).$$

Letting  $n \rightarrow +\infty$  and using boundedness condition from (4.16), we get

$$(\eta M_\tau h) \cdot h(x) = M_\tau h(x).$$

Hence  $h$  is an eigenvector of  $M_\tau$  associated to the eigenvalue  $\eta M_\tau h$ . Moreover, by the semigroup property,  $\eta M_{k\tau} h = (\eta M_\tau h)^k$ . Then, for all  $k \geq 0$ ,  $\eta M_{k\tau} h = e^{\lambda k \tau}$ , with  $\lambda = \tau^{-1} \log(\eta M_\tau h)$ . Next, for  $t \geq 0$  we have

$$M_\tau M_t h = e^{\lambda \tau} M_t h$$

so that  $M_t h$  is an eigenfunction of  $M_\tau$  with associated eigenvalue  $e^{\lambda \tau}$ . By uniqueness, there exists  $(c_t)_{t \geq 0}$  such that for all  $t \geq 0$ ,  $M_t h = c_t h$ . Moreover, by the semigroup property,  $c_{t+s} = c_t c_s$  for all  $t, s \geq 0$  and  $t \mapsto c_t$  is locally bounded using (4.15) and  $t \mapsto \|M_t V\|_{\mathcal{B}(V)}$  is locally bounded. Therefore, there exists  $\tilde{\lambda} \in \mathbb{R}$  such that  $c_t = e^{\tilde{\lambda} t}$  and as  $c_\tau = e^{\lambda \tau}$ , we finally get  $\tilde{\lambda} = \lambda$ . Then, for all  $t \geq 0$ ,  $M_t h = e^{\lambda t} h$ .

Let us proceed to the last step and provide a lower bound for  $h$ . Combining

$$M_{n\tau}\psi \geq M_{k\tau}(\mathbf{1}_K M_{(n-k)\tau}\psi) = M_{k\tau}(\mathbf{1}_K M_\tau((M_{(n-k-1)\tau}\psi/\psi)\psi))$$

for all  $k \leq n$  with (A3) and (A2), we get

$$M_{n\tau}\psi \geq c\beta(\eta M_{(n-k-1)\tau}\psi)M_{k\tau}(\mathbf{1}_K\psi).$$

Recalling (4.7) and dividing by  $\eta M_{n\tau}\psi$ , we obtain

$$\frac{M_{n\tau}\psi}{\eta M_{n\tau}\psi} \geq c\beta c_{k-1} (\beta^k \psi - \alpha^{k-1} (\alpha + \theta) V) \frac{\eta M_{(n-k-1)\tau}\psi}{\eta M_{n\tau}\psi}.$$

Let  $n \rightarrow \infty$ , the left-hand side converges to  $h$  and using from Lemma 4.1 *iii*) that  $\eta M_{(n-k-1)\tau}\psi/\eta M_{n\tau}\psi \geq 1/\Xi^{k+1}$ , and recalling the expression of  $c_k$  in (4.2), we obtain  $h \geq c(cr)^k (\psi - \alpha^{k-1}(\alpha + \theta)V/\beta)$ .

Considering now  $k = k(x) = \left\lfloor \log \left( \frac{\psi(x)}{V(x)} \frac{\beta}{2(\alpha + \theta)} \right) / \log(\mathbf{a}) \right\rfloor + 2$  and recalling that  $r = \beta^2/\Xi^2$  yields  $\psi - \alpha^{k-1} \frac{\alpha + \theta}{\beta} V \geq \psi/2$ . We get

$$h \geq c_1 \left( \frac{\psi}{V} \right)^q \psi, \quad \text{with } c_1 = c(cr)^{2 + \log(\frac{\beta}{2(\alpha + \theta)})/\log(\mathbf{a})} / 2 > 0$$

and  $q = \log(cr)/\log(\mathbf{a}) > 0$ , which ends the proof.  $\square$

**Remark 4.7.** Notice that the eigenfunction  $h$  built in this proof satisfies  $\nu(h/\psi) = 1$  and the constants in  $(V/\psi)^q \psi \lesssim h \lesssim V$  depend on this normalization. If we normalize  $h$  such that  $\|h\|_{\mathcal{B}(V)} = 1$  as in Theorem 2.1 we get  $c_1 d_2 (\psi/V)^q \psi \leq h \leq V$ .

We consider now the left eigenelement.

*Proof of Lemma 3.5.* Let us use again  $\eta = \nu(\cdot/\psi)$ . Applying Lemma 4.6 to  $\mu_1 = \eta$  and  $\mu_2 = \eta M_{n\tau}$  and using again (4.20), we get for  $k, n \geq 0$ ,

$$\|Q_{(k+n)\tau}^\eta - Q_{k\tau}^\eta\|_{\mathcal{M}(1+\kappa V_0)} \leq C'_2 \rho^{\lfloor k/p \rfloor} \left( \nu \left( \frac{V}{\psi} \right) + [\eta M_{n\tau}] \right),$$

where  $C'_2$  is defined in (4.21). Then, using Lemma 4.1 *ii*),  $V \leq R\psi$  on  $K$  and that  $\nu(K) = 1$ , we have

$$\|Q_{(k+n)\tau}^\eta - Q_{k\tau}^\eta\|_{\mathcal{M}(1+\kappa V_0)} \leq C'_2 \rho^{\lfloor k/p \rfloor} (R + \mathbf{a}^n R + \Theta).$$

Therefore, the sequence of probabilities  $(Q_{k\tau}^\eta)_{k \geq 0}$  satisfies the Cauchy criterion in  $\mathcal{M}(1 + \kappa V_0)$  and it then converges to a probability measure  $\pi \in \mathcal{M}(1 + \kappa V_0)$ . Similarly, applying Lemma 4.6 to  $\mu_1 = \mu$  and  $\mu_2 = \eta M_{n\tau}$ , we also have

$$\|Q_{(k+n)\tau}^\eta - Q_{k\tau}^\mu\|_{\mathcal{M}(1+\kappa V_0)} \leq C'_2 \rho^{\lfloor k/p \rfloor} ([\mu] + \mathbf{a}^n R + \Theta)$$

for any  $\mu \in \mathcal{M}(1 + \kappa V_0)$ . Letting  $n$  tend to infinity yields

$$\|\pi - Q_{k\tau}^\mu\|_{\mathcal{M}(1+\kappa V_0)} \leq C'_2([\mu] + \Theta) \rho^{\lfloor k/\mathfrak{p} \rfloor}. \quad (4.23)$$

Besides,  $\pi(h/\psi) \leq \pi(V/\psi) = \pi(V_0) < +\infty$  and we can define  $\gamma \in \mathcal{M}(\psi + \kappa V)$  by

$$\gamma(f) = \frac{\pi(f/\psi)}{\pi(h/\psi)}$$

for  $f \in \mathcal{B}(\psi + \kappa V) = \mathcal{B}(V)$ . Observe that  $\gamma(h) = 1$ . Next,

$$Q_{(k+1)\tau}^\eta(f/\psi) = Q_{k\tau}^\eta(M_\tau f/\psi) \frac{\eta M_{k\tau} \psi}{\eta M_{(k+1)\tau} \psi}. \quad (4.24)$$

Applying (3.6) to  $\mu = \eta M_\tau$ ,

$$\frac{\eta M_{k\tau} \psi}{\eta M_{(k+1)\tau} \psi} \xrightarrow{k \rightarrow +\infty} e^{-\lambda\tau}.$$

Then, letting  $k \rightarrow \infty$  in (4.24), we obtain  $\pi(f/\psi) = \pi(M_\tau f/\psi)e^{-\lambda\tau}$ , which ensures that  $\gamma$  is an eigenvector for  $M_\tau$ . As in the proof of Lemma 3.4, using the semigroup property and that  $t \mapsto \|M_t V\|_{\mathcal{B}(V)}$  is locally bounded, we get that for all  $t \in \mathbb{R}_+$  that  $\gamma M_t = e^{\lambda t} \gamma$ . Adding that  $\pi(f) = \gamma(f\psi)/\gamma(\psi)$  since  $\pi$  is a probability measure, (3.7) follows from (4.23).  $\square$

## 4.4 Proofs of Section 3.3

We start by proving a discrete version of Theorem 3.6.

**Proposition 4.8.** *Under Assumption A, there exists  $C > 0$  such that for all  $\mu \in \mathcal{M}_+(V)$  and all integers  $k \geq 0$  we have*

$$\|\mu M_{k\tau} - e^{\lambda k\tau} \mu(h)\gamma\|_{\mathcal{M}(V)} \leq C[\mu] e^{-\sigma k\tau} \min\{\mu M_{k\tau} \psi, \mu(V)e^{\lambda k\tau}\}, \quad (4.25)$$

where

$$\sigma = \frac{-\log \rho}{\mathfrak{p}\tau} > 0.$$

*Proof.* Using that

$$\|\pi - Q_{k\tau}^\mu\|_{\mathcal{M}(1+\kappa V_0)} = \sup_{f \in \mathcal{B}(1+\kappa V_0)} \left| \frac{\gamma(f\psi)}{\gamma(\psi)} - \frac{\mu M_{k\tau}(f\psi)}{\mu M_{k\tau} \psi} \right| = \left\| \frac{\gamma}{\gamma(\psi)} - \frac{\mu M_{k\tau}}{\mu M_{k\tau} \psi} \right\|_{\mathcal{M}(\psi + \kappa V)}$$

and multiplying (4.23) by  $\mu M_{k\tau} \psi$ , we get

$$\left\| \mu M_{k\tau} \psi \frac{\gamma}{\gamma(\psi)} - \mu M_{k\tau} \right\|_{\mathcal{M}(\psi + \kappa V)} \leq C'_2 \rho^{\lfloor k/\mathfrak{p} \rfloor} ([\mu] + \Theta) \mu M_{k\tau} \psi. \quad (4.26)$$

Moreover,  $h \in \mathcal{M}(\psi + \kappa V)$  since  $h \lesssim V$ . As  $\gamma(h) = 1$ , the previous inequality applied to the eigenfunction  $h$  yields

$$\left| \frac{\mu M_{k\tau} \psi}{\gamma(\psi)} - \mu(h) e^{\lambda k \tau} \right| \leq C'_2 \rho^{\lfloor k/p \rfloor} ([\mu] + \Theta) \mu M_{k\tau} \psi.$$

Then, recalling that  $\psi \leq V$ , we have

$$\begin{aligned} \left\| \mu M_{k\tau} \psi \frac{\gamma}{\gamma(\psi)} - e^{\lambda k \tau} \mu(h) \gamma \right\|_{\mathcal{M}(\psi + \kappa V)} &= \left| \frac{\mu M_{k\tau} \psi}{\gamma(\psi)} - e^{\lambda k \tau} \mu(h) \right| \|\gamma\|_{\mathcal{M}(\psi + \kappa V)} \\ &\leq C'_2 \rho^{\lfloor k/p \rfloor} ([\mu] + \Theta) \mu M_{k\tau} \psi \times (1 + \kappa) \gamma(V). \end{aligned} \quad (4.27)$$

Combining (4.26) and (4.27), by triangular inequality, we get

$$\left\| \mu M_{k\tau} - \gamma e^{\lambda k \tau} \mu(h) \right\|_{\mathcal{M}(\psi + \kappa V)} \leq C'_2 \rho^{\lfloor k/p \rfloor} ([\mu] + \Theta) \mu M_{k\tau} \psi (1 + (1 + \kappa) \gamma(V)). \quad (4.28)$$

This gives the first part of (4.25). Finally, by integration of (4.16),

$$\mu M_{k\tau} \psi \leq d_1^{-1} (\alpha + \theta) \nu(M_{k\tau} \psi / \psi) \mu(V). \quad (4.29)$$

Adding that  $[\gamma] \leq \Theta$  according to Lemma 4.1 *ii*) and  $\gamma M_{k\tau} = e^{\lambda k \tau} \gamma$  from Lemma 3.5, Lemma 4.5 applied to  $\mu = \gamma$  yields  $\nu(M_{k\tau} \psi / \psi) \leq C_1 e^{\lambda k \tau}$  for  $k \geq p$  and we obtain  $\mu M_{k\tau} \psi \leq C_1 d_1^{-1} (\alpha + \theta) \mu(V) e^{\lambda k \tau}$ . It proves (4.25) for  $k \geq p$  with

$$C = C'_2 (1 + \Theta) (1 + (1 + \kappa) \gamma(V)) \max(1, C_1 d_1^{-1} (\alpha + \theta)).$$

The fact that (4.25) holds for some constant  $C$  also for  $k \leq p$  comes directly from (4.28), (4.29), Lemma 4.1 *iii*) and  $\nu(K) = 1$ .  $\square$

Next, we extend this result to continuous time.

*Proof of Theorem 3.6.* We introduce the following constant

$$C_0 = \sup_{s \leq \tau} \max \left\{ \left\| \frac{M_s V}{V} \right\|_{\infty}, \left\| \frac{\psi}{M_s \psi} \right\|_{\infty} \right\},$$

which is finite under Assumption (3.9). Let  $t \geq 0$  and  $(k, \varepsilon) \in \mathbb{N} \times [0, \tau)$  be such that  $t = k\tau + \varepsilon$ . Using that  $\mu M_t h = e^{\lambda t} \mu(h)$ , we have

$$\left\| \mu M_t - e^{\lambda t} \mu(h) \gamma \right\|_{\mathcal{M}(V)} = \left\| \mu M_{k\tau + \varepsilon} - \mu(M_{k\tau + \varepsilon} h) \gamma \right\|_{\mathcal{M}(V)}.$$

Then, from (4.25) with  $\mu := \mu M_\varepsilon$ , there exists  $C > 0$  such that for all integers  $k \geq 0$ ,

$$\begin{aligned} \left\| \mu M_t - e^{\lambda t} \mu(h) \gamma \right\|_{\mathcal{M}(V)} &\leq C [\mu M_\varepsilon] e^{-\sigma k \tau} \min \{ \mu M_{k\tau + \varepsilon} \psi, \mu M_\varepsilon V e^{\lambda k \tau} \} \\ &\leq C C_0^2 e^{\sigma \tau} \max \{ 1, C_0 e^{|\lambda| \tau} \} [\mu] e^{-\sigma t} \min \{ \mu M_t \psi, \mu(V) e^{\lambda t} \}, \end{aligned}$$

which ends the proof.  $\square$

Now we turn to the proof of Corollary 3.7.

*Proof of Corollary 3.7.* For convenience and without loss of generality, we assume that  $V \geq \mathbf{1}$ . Then,  $\gamma(\mathbf{1}) < \infty$ . Next, if  $\gamma(\mathbf{1}) = 0$ , then  $\gamma(\mathcal{X}) = 0$ . In this case,  $\gamma = 0$ , which is absurd because  $\gamma(\psi) > 0$  and  $\psi > 0$ . Therefore,  $\gamma(\mathbf{1}) > 0$ .

We set  $\pi(\cdot) = \gamma(\cdot)/\gamma(\mathbf{1})$  and we have by triangular inequality

$$\begin{aligned} \left\| \frac{\mu M_t}{\mu M_t \mathbf{1}} - \pi \right\|_{\text{TV}} &\leq \frac{e^{\lambda t}}{\mu M_t \mathbf{1}} \left\| e^{-\lambda t} \mu M_t - \pi e^{-\lambda t} \mu M_t \mathbf{1} \right\|_{\mathcal{M}(V)} \\ &\leq \frac{e^{\lambda t}}{\mu M_t \mathbf{1}} \left( \left\| e^{-\lambda t} \mu M_t - \gamma \mu(h) \right\|_{\mathcal{M}(V)} + |\gamma(\mathbf{1})\mu(h) - e^{-\lambda t} \mu M_t \mathbf{1}| \pi(V) \right). \end{aligned}$$

From (2.2), we have  $\left\| e^{-\lambda t} \mu M_t - \gamma \mu(h) \right\|_{\mathcal{M}(V)} \leq C\mu(V)e^{-\omega t}$ . Using this estimate with  $V \geq \mathbf{1}$ , we also have

$$|\gamma(\mathbf{1})\mu(h) - e^{-\lambda t} \mu M_t \mathbf{1}| \leq C\mu(V)e^{-\omega t}. \quad (4.30)$$

Combining the three last estimates yields

$$\left\| \frac{\mu M_t}{\mu M_t \mathbf{1}} - \pi \right\|_{\text{TV}} \leq C \frac{e^{\lambda t}}{\mu M_t \mathbf{1}} \mu(V) e^{-\omega t} (1 + \pi(V)). \quad (4.31)$$

Now on the first hand, Equation (4.30) also gives  $e^{-\lambda t} \mu M_t \mathbf{1} \geq \gamma(\mathbf{1})\mu(h) - C\mu(V)e^{-\omega t}$ , and for any  $t \geq t(\mu) = \frac{1}{\omega} \log \left( \frac{2C}{\gamma(\mathbf{1})} \frac{\mu(V)}{\mu(h)} \right)$ , we have

$$e^{-\lambda t} \mu M_t \mathbf{1} \geq \mu(h)\gamma(\mathbf{1})/2. \quad (4.32)$$

Plugging (4.32) in (4.31) yields (3.10) when  $t \geq t(\mu)$ . Otherwise,

$$\left\| \frac{\mu M_t}{\mu M_t \mathbf{1}} - \pi \right\|_{\text{TV}} \leq 1 \leq e^{-\omega t} e^{\omega t(\mu)} \leq C \frac{\mu(V)}{\mu(h)} e^{-\omega t},$$

which ends the proof.  $\square$

## 4.5 Proofs of Section 2

*Proof of Theorem 2.1 i).* We assume that Assumption **A** is satisfied by  $(V, \psi)$  for a set  $K$ , constants  $\alpha, \beta, \theta, c, d$  and a probability  $\nu$ . Then, from Lemmas 3.4 and 3.5, there exist eigenelements  $(\gamma, h, \lambda)$  such that

$$\beta \leq e^{\lambda \tau} \leq \alpha + \theta, \quad c_1 d_2 (\psi/V)^q \psi \leq h \leq V.$$

We check now that  $(V, h)$  satisfies also Assumption **A** with the same set  $K$  and constant  $\alpha$  as  $(V, \psi)$  but other constants  $\beta', \theta', c', d'$  and an other probability measure  $\nu'$ . First the inequality  $V/h \leq \frac{1}{c_1 d_2} (V/\psi)^{q+1}$  ensures that  $\sup_K V/h < \infty$  since  $\sup_K V/\psi < \infty$ .

$\infty$ . Moreover  $h$  satisfies (A2) with  $\beta' = \exp(\lambda\tau) \geq \beta > \alpha$ . Recalling that  $R = \sup_K V/\psi < \infty$ , we also have

$$\sup_K \frac{\psi}{h} \leq \frac{R^q}{c_1 d_2}.$$

Adding that  $(V, \psi)$  satisfies (A1) with constants  $\alpha, \theta$  yields  $M_\tau V \leq \alpha V + \theta' \mathbf{1}_K h$  with  $\theta' = \theta \frac{R^q}{c_1 d_2}$ , which gives (A1) for  $(V, h)$ . We use now (A3) and (A2) for  $(V, \psi)$  and get for  $x \in K$

$$\delta_x M_\tau(fh) \geq c\nu \left( \frac{fh}{\psi} \right) \delta_x M_\tau \psi \geq c\beta\nu \left( \frac{fh}{\psi} \right) \psi(x).$$

Using again that  $M_\tau h = e^{\lambda\tau} h$ , we obtain for  $x \in K$ ,  $\delta_x M_\tau(fh) \geq c'\nu'(f)\delta_x M_\tau h$ , with

$$\nu' = \frac{\nu(\cdot \frac{h}{\psi})}{\nu(\frac{h}{\psi})}, \quad c' = c\beta e^{-\lambda\tau} \nu \left( \frac{h}{\psi} \right) \inf_K \frac{\psi}{h} \geq \frac{\beta}{\alpha + \theta} \frac{c_1 d_2}{R^{q+1}} > 0.$$

Finally, (A4) is satisfied since

$$\sup_{x \in K} \frac{M_{n\tau} h(x)}{h(x)} = e^{\lambda n\tau} = \nu' \left( \frac{M_{n\tau} h}{h} \right).$$

Therefore, every result stated above holds replacing  $\psi$  by  $h$  and the constants  $\beta, \theta, c, d$  of Assumption **A** by  $\beta', \theta', c', d'$  defined above. In particular, for all  $\mu \in \mathcal{M}_+(V)$ , (3.7) becomes

$$\left\| \gamma(\cdot h) - \frac{\mu M_{k\tau}(\cdot h)}{e^{\lambda k\tau} \mu(h)} \right\|_{\mathcal{M}(1+\kappa V/h)} \leq C \left( \frac{\mu(V)}{\mu(h)} + \Theta' \right) \rho^{\lfloor k/p \rfloor},$$

where  $\Theta' = \theta' / (\beta' - \alpha)$  and

$$C = \max \{ 2\mathbf{a}^{-p} + \kappa C_1 (1 + 2\Theta \mathbf{a}^{-p}), 2(1 + \kappa \Theta) \mathbf{a}^{-(p+p)} + \kappa \} \max \left( 1, \frac{(\alpha + \theta')^p}{\beta^p} \right)$$

and  $\rho$  is defined in (3.5),  $\mathbf{p}$  in (3.4),  $\mathbf{a}$  in (3.3),  $\kappa$  in Proposition 3.3,  $\Theta$  in (4.1), and  $p, C_1$  in (4.4) (replacing  $\beta, \theta, c, d$  by  $\beta', \theta', c', d'$  in each definition). Using that

$$\left\| \gamma(\cdot h) - \frac{\mu M_{k\tau}(\cdot h)}{e^{\lambda k\tau} \mu(h)} \right\|_{\mathcal{M}(1+\kappa V/h)} = \left\| \gamma - \frac{\mu M_{k\tau}}{e^{\lambda k\tau} \mu(h)} \right\|_{\mathcal{M}(h+\kappa V)} \geq \kappa \left\| \gamma - \frac{\mu M_{k\tau}}{e^{\lambda k\tau} \mu(h)} \right\|_{\mathcal{M}(V)},$$

multiplying by  $e^{\lambda k\tau} \mu(h)$  and using that  $h \leq V$ , we get

$$\left\| e^{\lambda k\tau} \mu(h) \gamma - \mu M_{k\tau} \right\|_{\mathcal{M}(V)} \leq \frac{C}{\kappa \rho} \mu(V) (1 + \Theta') e^{-\omega k\tau} e^{\lambda k\tau},$$

where  $\omega = -\log \rho / \mathbf{p}\tau$ . Let  $t \geq 0$  and  $(k, \varepsilon) \in \mathbb{N} \times [0, \tau)$  be such that  $t = k\tau + \varepsilon$ .

Applying the above inequality to the measure  $\mu M_\varepsilon$  in place of  $\mu$  yields

$$\left\| e^{\lambda t} \mu(h) \gamma - \mu M_t \right\|_{\mathcal{M}(V)} \leq \frac{C}{\kappa \rho} \mu(M_\varepsilon V) (1 + \Theta') e^{-\omega k\tau} e^{\lambda k\tau} \leq C' \mu(V) e^{-\omega t},$$

where  $C' = \frac{C}{\kappa \rho} (1 + \Theta') e^{\omega\tau} e^{|\lambda|\tau} \sup_{s \leq \tau} \left\| \frac{M_s V}{V} \right\|_\infty$ . Adding that uniqueness is a direct consequence of  $\omega > 0$  ends the proof of Theorem 2.1 *i*).  $\square$

*Proof of Theorem 2.1 ii).* The proof follows the usual equivalence in Harris Theorem [36, Chapter 15]. It also used the necessary condition of small set obtained in [21, Theorem 2.1]. Assume that there exist a positive measurable function  $V$ , a triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$ , and constants  $C, \omega > 0$  such that (2.1) and (2.2) hold. Without loss of generality we can suppose that  $\|h\|_{\mathcal{B}(V)} = \gamma(h) = 1$ . It remains to check that  $(V, h)$  satisfies Assumption **A**.

Fix  $R > \gamma(V)$  and  $\tau > 0$  such that

$$e^{-\omega\tau/2}C(R + \gamma(V)) < 1 - \frac{\gamma(V)}{R}. \quad (4.33)$$

It ensures that

$$\alpha := e^{\lambda\tau} \left( Ce^{-\omega\tau} + \frac{\gamma(V)}{R} \right) < \beta := e^{\lambda\tau}.$$

Using (2.1), we obtain that (A2) and (A4) are satisfied by  $h$  with  $d = 1$  and any probability measure  $\nu$ , which ends the proof.

By (2.2), we have for all  $x \in \mathcal{X}$ ,  $e^{-\lambda t}M_t V(x) - h(x)\gamma(V) \leq CV(x)e^{-\omega t}$ . We define  $K = \{x \in \mathcal{X}, V(x) \leq Rh(x)\}$ , which is not empty since  $\|h\|_{\mathcal{B}(V)} = 1$  and  $R > \gamma(V) \geq \gamma(h) = 1$ . Writing  $\theta = \gamma(V)e^{\lambda\tau}$  and using  $h(x) = \mathbf{1}_{K^c}(h(x)/V(x))V(x) + \mathbf{1}_K h(x)$ , we get

$$M_\tau V(x) \leq \alpha V(x) + \mathbf{1}_K \theta h(x)$$

for all  $x \in \mathcal{X}$ . Therefore, (A1) holds for  $(V, h)$  and it remains to prove (A3).

We define the probability measure  $\pi := \gamma(\cdot h)$  and we use the Hahn-Jordan decomposition of the following family of signed measure indexed by  $x \in \mathcal{X}$ ,

$$\nu^x = \frac{\delta_x M_{\tau/2}(\cdot h)}{e^{\lambda\tau/2}h(x)} - \pi = \nu_+^x - \nu_-^x.$$

As  $h \leq V$ , Equation (2.2) with  $t = \tau/2$  and  $\mu = \delta_x$  yields

$$\nu_\pm^x(\mathbf{1}) \leq \nu_\pm^x(V/h) \leq \|\nu^x\|_{\mathcal{M}(V/h)} = \left\| \frac{\delta_x M_{\tau/2}}{e^{\lambda\tau/2}h(x)} - \gamma \right\|_{\mathcal{M}(V)} \leq C \frac{V(x)}{h(x)} e^{-\omega\tau/2}. \quad (4.34)$$

For every  $f \in \mathcal{B}_+(V/h)$  and  $x \in \mathcal{X}$  we have

$$\frac{\delta_x M_\tau(hf)}{e^{\lambda\tau}h(x)} = \frac{\delta_x M_{\tau/2}}{e^{\lambda\tau/2}h(x)} \left( \frac{M_{\tau/2}(hf)}{e^{\lambda\tau/2}h} h \right) \geq (\pi - \nu_-^x) \left( \frac{M_{\tau/2}(hf)}{e^{\lambda\tau/2}h} \right). \quad (4.35)$$

Next,

$$\pi \left( \frac{M_{\tau/2}(hf)}{e^{\lambda\tau/2}h} \right) = \frac{\gamma M_{\tau/2}(hf)}{e^{\lambda\tau/2}} = \frac{e^{\lambda\tau/2}\gamma(hf)}{e^{\lambda\tau/2}} = \pi(f) \quad (4.36)$$

and writing  $(\nu^y(f))_+$  the positive part of the real number  $\nu^y(f)$ ,

$$\nu_-^x \left( \frac{M_{\tau/2}(hf)}{e^{\lambda\tau/2}h} \right) = \int_{\mathcal{X}} \frac{\delta_y M_{\tau/2}(hf)}{e^{\lambda\tau/2}h(y)} \nu_-^x(dy) \leq \int_{\mathcal{X}} (\pi(f) + (\nu^y(f))_+) \nu_-^x(dy). \quad (4.37)$$

Combining (4.35) with (4.36) and (4.37), we get

$$\frac{\delta_x M_{\tau}(hf)}{e^{\lambda\tau}h(x)} \geq \pi(f)(1 - \nu_-^x(\mathbf{1})) - \int_{\mathcal{X}} (\nu^y(f))_+ \nu_-^x(dy).$$

The minimality property of the Hahn-Jordan decomposition entails that  $\nu_-^x \leq \pi$  and (4.34) yields  $\nu_-^x(\mathbf{1}) \leq CR e^{-\omega\tau/2}$  for  $x \in K$ . We deduce that for all  $x \in K$ ,

$$\frac{\delta_x M_{\tau}(hf)}{e^{\lambda\tau}h(x)} \geq \pi(f)(1 - CR e^{-\omega\tau/2}) - \int (\nu^y(f))_+ \pi(dy) =: \eta(f).$$

We consider now the infimum measure  $\nu_0$  of the left hand side. More precisely, using [21, Lemma 5.2] we can define the mesure  $\nu_0$  by

$$\nu_0(A) = \inf \left\{ \sum_{i=1}^n \frac{\delta_{x_i} M_{\tau}(h\mathbf{1}_{B_i})}{e^{\lambda\tau}h(x_i)} : n \geq 1; (x_1, \dots, x_n) \in \mathcal{X}^n; (B_1, \dots, B_n) \in \mathfrak{P}(A) \right\},$$

where  $A$  is a measurable subset of  $\mathcal{X}$  and  $\mathfrak{P}(A)$  is the set of finite partitions of  $A$  formed by measurable sets of  $\mathcal{X}$ . Besides for any measurable partition  $B_1, \dots, B_n$  of  $\mathcal{X}$ , we have  $\sum_{i=1}^n (\nu^y(\mathbf{1}_{B_i}))_+ \leq \nu_+^y(\mathcal{X}) \leq C e^{-\omega\tau/2} V(y)/h(y)$  from (4.34). Then

$$\sum_{i=1}^n \eta(\mathbf{1}_{B_i}) \geq 1 - CR e^{-\omega\tau/2} - \int_{\mathcal{X}} \sum_{i=1}^n (\nu^y(\mathbf{1}_{B_i}))_+ \pi(dy) \geq c,$$

where  $c = 1 - C e^{-\omega\tau/2}(R + \gamma(V)) > 0$  using (4.33). Adding that

$$\eta(\mathbf{1}_{K^c}) \leq \eta \left( \frac{V}{R\psi} \right) \leq \pi \left( \frac{V}{R\psi} \right) = \frac{\gamma(V)}{R},$$

we obtain that  $\nu_0(\mathcal{X}) \geq c$  and  $\nu_0(K^c) < c$ . As a conclusion,  $\nu_0$  is a positive measure such that  $\nu_0(K) > 0$  and for any  $x \in \mathcal{X}$  and  $f \in \mathcal{B}_+(V/h)$ ,

$$\frac{\delta_x M_{\tau}(hf)}{e^{\lambda\tau}h(x)} \geq \nu_0(f)$$

which yields (A3) and ends the proof.  $\square$

We end this section by proving the sufficient conditions of Section 2.2.

*Proof of Proposition 2.2.* Let  $C > 0$  be such that  $C^{-1}\psi \leq \varphi \leq C\psi$ . By assumptions  $\mathcal{L}\psi \geq b\psi$  and  $\mathcal{L}\varphi \leq \xi\varphi$  so that we have for all  $t \geq 0$

$$M_t\psi \geq \psi + b \int_0^t M_s\psi ds \quad \text{and} \quad M_t\varphi \leq \varphi + \xi \int_0^t M_s\varphi ds,$$



which yields by Grönwall's lemma  $M_t\psi \geq e^{bt}\psi$  and  $M_t\psi \leq CM_t\varphi \leq Ce^{\xi t}\varphi \leq C^2e^{\xi t}\psi$ . Similarly, setting  $\phi = V - \frac{\zeta}{b-a}\psi$ , we have  $\mathcal{L}\phi \leq aV + \zeta\psi - \frac{b\zeta}{b-a}\psi = a\phi$ , so by Grönwall's lemma  $M_t\phi \leq e^{at}\phi$  and

$$M_tV \leq e^{at}\left(V - \frac{\zeta}{b-a}\psi\right) + \frac{\zeta}{b-a}M_t\psi \leq e^{at}V + \frac{C^2\zeta}{b-a}e^{\xi t}\psi.$$

Consider now  $\tau, R > 0$ , and

$$\alpha = e^{a\tau} + \frac{C^2\zeta}{(b-a)R}e^{\xi\tau}, \quad \beta = e^{b\tau}, \quad \theta = \frac{C^2\zeta}{b-a}e^{\xi\tau}, \quad R_0 = \frac{\theta}{e^{b\tau} - e^{a\tau}}$$

positive constants. Defining  $K = \{x \in \mathcal{X}, V(x) \leq R\psi(x)\}$ , we get  $\psi \leq V/R$  on  $K^c$  and  $M_\tau V \leq \alpha V + \theta \mathbf{1}_K \psi$ . Besides for  $R > R_0$ ,  $\alpha < \beta$ . So Assumptions (A1)-(A2) are verified for  $K = \{V \leq R\psi\}$ ,  $R > R_0$ , and constants  $\alpha, \beta, \theta$  defined just above.  $\square$

*Proof of Proposition 2.3.* Let  $\psi : \mathcal{X} \rightarrow (0, \infty)$  and define  $\nu = (\#K)^{-1} \sum_{x \in K} \delta_x$  the uniform measure on  $K$ , where  $\#K$  stands for the cardinal of  $K$ . We have for all  $f \geq 0$  and  $x, y \in K$ ,

$$\delta_x M_\tau(f\psi) \geq \delta_x M_\tau(\{y\})f(y)\psi(y) \geq cf(y)M_\tau\psi(x).$$

where

$$c = \min_{x, y \in K} \frac{\psi(y)\delta_x M_\tau(\{y\})}{M_\tau\psi(x)} > 0$$

using the irreducibility condition  $\delta_x M_\tau(\{y\}) > 0$ . Integrating with respect to  $\nu$  shows that (A3) holds and Assumption (A4) is trivially satisfied with  $d = 1/\#K$ .  $\square$

## 5 Applications

### 5.1 Convergence to quasi-stationary distribution

Let  $(X_t)_{t \geq 0}$  be a càd-làg Markov process on the state space  $\mathcal{X} \cup \{\partial\}$ , where  $\mathcal{X}$  is measurable space and  $\partial$  is an absorbing state. In this section, we apply the results to the (non-conservative) semigroup defined for any measurable bounded function  $f$  on  $\mathcal{X}$  and any  $x \in \mathcal{X}$  by

$$M_t f(x) = \mathbb{E}_x [f(X_t) \mathbf{1}_{X_t \neq \partial}].$$

We assume that there exists a positive function  $V$  such that for any  $t > 0$ , there exists  $C_t > 0$  such that,  $\mathbb{E}_x[V(X_t)] \leq C_t V(x)$  for any  $x \in \mathcal{X}$ . This ensures that the

semigroup  $M$  acts on  $\mathcal{B}(V)$  and that we can use the framework of Section 2. A quasi-stationary distribution (QSD) is a probability law  $\pi$  on  $\mathcal{X}$  such that

$$\forall t \geq 0, \mathbb{P}_\pi(X_t \in \cdot | X_t \neq \partial) = \pi(\cdot).$$

Corollary 3.7 directly gives existence and uniqueness of a QSD and quantitative estimates for the convergence. Recall that  $\mathcal{P}(V)$  stands for the set of probability measures which integrate  $V$ .

**Theorem 5.1.** *Assume that  $(M_t)_{t \geq 0}$  satisfies Assumption **A** with  $\inf_{\mathcal{X}} V > 0$ . Then, there exist a unique quasi-stationary distribution  $\pi \in \mathcal{P}(V)$ , and  $\lambda_0 > 0, h \in \mathcal{B}_+(V)$ ,  $C, w > 0$  such that for all  $\mu \in \mathcal{P}(V)$  and  $t \geq 0$*

$$\|e^{\lambda_0 t} \mathbb{P}_\mu(X_t \in \cdot) - \mu(h)\pi\|_{\text{TV}} \leq C\mu(V)e^{-\omega t},$$

and

$$\|\mathbb{P}_\mu(X_t \in \cdot | X_t \neq \partial) - \pi\|_{\text{TV}} \leq C \frac{\mu(V)}{\mu(h)} e^{-\omega t}.$$

It extends and complements recent results, see *e.g.* [22] for various interesting examples and discussions below for comparisons of statements. In particular, it relaxes the conditions of boundedness for  $\psi$  and provides a quantitative estimate for exponential convergence of the conditional distribution to the QSD using the eigenfunction  $h$ .

As an application, we consider the simple but interesting case of a continuous time random walk on integers, with jumps  $+1$  and  $-1$ , absorbed at 0. We obtain optimal results for the exponential convergence to the QSD. Let us consider the Markov process  $X$  whose transition rates and generator are given by the linear operator defined for any  $n \in \mathbb{N}$  and  $f : \mathbb{N} \rightarrow \mathbb{R}$  by

$$\mathcal{L}f(n) = b_n(f(n+1) - f(n)) + d_n(f(n-1) - f(n)),$$

where  $b_i = b > 0$ ,  $d_i = d > 0$  for any  $i \geq 2$ ,  $b_1, d_1 > 0$ ,  $b_0 = d_0 = 0$ . This process is a Birth and Death (BD) process which follows a simple random walk before reaching 1. If  $d \geq b$ , this process is almost surely absorbed at 0. The convergence in law of such processes conditionally on non-absorption has been studied in many works [78, 77, 81, 63, 1, 46, 60, 85, 52].

The necessary and sufficient condition for  $\xi$ -positive recurrence of BD processes is known from the work of van Doorn [78]. More precisely here, the fact that there exists  $\lambda > 0$  such that for any  $x > 0$  and  $i > 0$ ,  $e^{-\lambda t} \mathbb{P}_x(X_t = i)$  converges to a positive finite limit as  $t \rightarrow \infty$  is given by the following condition

$$\text{(H)} \quad \Delta := (\sqrt{b} - \sqrt{d})^2 + b_1 \left( \sqrt{d/b} - 1 \right) - d_1 > 0.$$

We notice that  $b = d$  is excluded by condition **(H)** and indeed in this case  $t \mapsto \mathbb{P}(X_t \neq 0)$  decreases polynomially. Similarly, the case  $b_1 = b$  and  $d_1 = d$  is excluded and there is an additional linear term in the exponential decrease of  $\mathbb{P}_x(X_t = i)$ . Moreover we know from [77] that condition **(H)** ensures that  $\mathbb{P}_x(X_t \in \cdot | X_t \neq 0)$  converges to the unique QSD  $\pi$  for any  $x > 0$ . To the best of our knowledge, under Assumption **(H)**, the speed of convergence of  $e^{-\lambda t} \mathbb{P}_x(X_t = i)$  or  $\mathbb{P}_x(X_t = i | X_t \neq 0)$  and the extension of the convergence to infinite support masses were not proved, see *e.g.* [77, page 695]. For a subset of parameters satisfying **(H)**, [81] obtains the convergence to QSDs for non-compactly supported initial measures  $\mu$  such that  $\mu(V) < \infty$ . We obtain below quantitative exponential estimates for the full range of parameters given by **(H)** and non-compactly supported initial measures.

More precisely, we set  $\mathcal{X} = \mathbb{N} \setminus \{0\} = \{1, \dots\}$ ,  $V : n \mapsto \sqrt{d/b}^n$ ,  $\psi : n \mapsto \eta^n$ , for  $n \in \mathcal{X}$ , where  $\eta = \sqrt{d/b} - \Delta/2b_1 \in (0, \sqrt{d/b})$ .

**Corollary 5.2.** *Under Assumption **(H)**, there exists a unique QSD  $\pi \in \mathcal{P}(V)$ , and  $\lambda_0 > 0, h \in \mathcal{B}_+(V)$  and  $C, w > 0$  such that for all  $\mu \in \mathcal{P}(V)$  and  $t \geq 0$ ,*

$$\|e^{\lambda_0 t} \mathbb{P}_\mu(X_t \in \cdot) - \mu(h)\pi\|_{\text{TV}} \leq C\mu(V)e^{-\omega t}$$

and

$$\|\mathbb{P}_\mu(X_t \in \cdot | X_t \neq 0) - \pi\|_{\text{TV}} \leq C \frac{\mu(V)}{\mu(h)} e^{-\omega t}.$$

Note that the constants above can be explicitly derived from Lemma 3.4, that these estimates hold for non-compactly supported initial laws and that  $V$  and  $\psi$  are not eigenelements. As perspectives, we expect this statement to be generalized to BD processes where  $b_n, d_n$  are constant outside some compact set of  $\mathbb{N}$ . Finally, we hope that the proof will help to study the non-exponential decrease of the non-absorption probability, in particular for random walks, corresponding to  $b = b_1, d = d_1$ .

*Proof of Corollary 5.2.* For  $u \geq 1$ , let  $\varphi_u : n \mapsto u^n$  for  $n \geq 1$  and  $\varphi_u(0) = 0$ . We have  $\mathcal{L}\varphi_u(n) = \lambda_u(n)\varphi_u(n)$ , for any  $n \in \mathbb{N}$ , where

$$\lambda_u(n) = \lambda_u = b(u-1) + d(1/u-1) \quad (n \geq 2), \quad \lambda_u(1) = b_1(u-1) - d_1.$$

We set

$$a = \inf_{u>0} \lambda_u = \lambda_{\sqrt{d/b}} = -(\sqrt{d} - \sqrt{b})^2, \quad \zeta = \Delta \frac{V(1)}{\psi(1)}.$$

Note that from **(H)**,  $\zeta > 0$ . Then, setting  $V(0) = \psi(0) = 0$ , we have on  $\mathbb{N} = \mathcal{X} \cup \{0\}$  that  $V = \varphi_{\sqrt{d/b}}$  and

$$\mathcal{L}V = aV + \zeta \mathbf{1}_{\{n=1\}}\psi \leq aV + \zeta\psi. \tag{5.1}$$

Moreover, on  $\mathbb{N}$ ,  $\psi = \varphi_\eta$  and

$$b\psi \leq \mathcal{L}\psi \leq \xi\psi \quad (5.2)$$

with  $b = \min(\lambda_\eta, \lambda_\eta(1)) = \min(\lambda_\eta, a + \frac{\Delta}{2}) > a = \inf_{u>0} \lambda_u$ , and  $\xi = \max(\lambda_\eta, \lambda_\eta(1))$ .

Using now a classical localization argument (see Appendix 6.2 for details), the drift conditions (5.1)-(5.2) ensure that for any  $n \geq 1$  and  $t \geq 0$ ,

$$\mathbb{E}_n[V(X_t)] \leq V(x) + \int_0^t \mathbb{E}_n[(aV + \zeta\psi)(X_s)]ds, \quad (5.3)$$

$$\psi(x) + \int_0^t \mathbb{E}_n[b\psi(X_s)]ds \leq \mathbb{E}_x[\psi(X_t)] \leq \psi(x) + \int_0^t \mathbb{E}_n[\xi\psi(X_s)]ds. \quad (5.4)$$

Considering the generator  $\mathcal{L}$  of the semigroup  $M_t f(x) = \mathbb{E}[f(X_t)1_{X_t \neq 0}]$  for  $x \in \mathcal{X}$  and  $f \in \mathcal{B}(V)$  and recalling the definition of Section 2.2, these inequalities ensure

$$\mathcal{L}V \leq aV + \zeta\psi, \quad \mathcal{L}\psi \geq b\psi, \quad \mathcal{L}\psi \leq \xi\psi.$$

Finally, the fact that  $b_i, d_i > 0$  for  $i \geq 1$  ensures  $\delta_i M_t(\{j\}) > 0$  for any  $i, j \in \mathcal{X}$  and  $t > 0$ . Then combining Propositions 2.2 and 2.3 ensures that Assumption **A** holds for  $M$  with the functions  $(V, \psi)$ . Applying Theorem 5.1 ends the proof.  $\square$

## 5.2 The growth-fragmentation equation

In this section we apply our general result to the growth-fragmentation partial differential equation

$$\partial_t u_t(x) + \partial_x u_t(x) + B(x)u_t(x) = \int_0^1 B\left(\frac{x}{z}\right)u_t\left(\frac{x}{z}\right)\frac{\wp(dz)}{z} \quad (5.5)$$

for  $t, x > 0$ . This nonlocal partial differential equation is complemented with the zero flux boundary condition  $u_t(0) = 0$  for all  $t > 0$  and an initial data  $u_0 = \mu$ . The unknown  $u_t(x)$  represents the population density at time  $t$  of some ‘‘particles’’ with ‘‘size’’  $x > 0$ , which can be for instance the size of a cell [32, 49], the length of a polymer fibril [40], or the window size in data transmission over the Internet [20, 10]. Each particle grows with speed 1, and splits with rate  $B(x)$  to produce smaller particles of sizes  $zx$  with  $0 < z < 1$  distributed with respect to the fragmentation kernel  $\wp$ .

We assume that  $B : [0, \infty) \rightarrow [0, \infty)$  is a continuously differentiable increasing function and  $\wp$  is a finite positive measure on  $[0, 1]$  for which there exist  $z_0 \in (0, 1)$ ,  $\epsilon \in [0, z_0]$  and  $c_0 > 0$  such that

$$\wp(dz) \geq \frac{c_0}{\epsilon} \mathbf{1}_{[z_0-\epsilon, z_0]}(z)dz \quad \text{if } \epsilon > 0 \quad \text{or} \quad \wp \geq c_0 \delta_{z_0} \quad \text{if } \epsilon = 0. \quad (5.6)$$

For any  $r \in \mathbb{R}$  we denote by  $\wp_r \in (0, +\infty]$  the moment of order  $r$  of  $\wp$

$$\wp_r = \int_0^1 z^r \wp(dz).$$

Notice that Assumption (5.6) implies that  $r \mapsto \wp_r$  is strictly decreasing. The conservation of mass during the fragmentation leads to impose

$$\wp_1 = 1.$$

The zero order moment  $\wp_0$  represents the mean number of fragments. The conditions above ensure that  $\wp_0 > 1$  and as a consequence the growth-fragmentation equation we consider is not conservative. The conservative form where  $\wp_1 = 1$  is replaced by  $\wp_0 = 1$  also appears in some situations [20, 10, 18]. In this case, the eigenelements are given by  $h(x) = 1$ ,  $\lambda = 0$ , and the classical theory of the conservative Harris theorem applies [17, 18]. Here we are interested in the more challenging non-conservative case.

An important feature in the long time behavior of the (non-conservative) growth-fragmentation equation is the property of asynchronous exponential growth [82], which refers to a separation of the variables  $t$  and  $x$  when time  $t$  becomes large: the size repartition of the population stabilizes and the total mass grows exponentially in time. This question attracted a lot of attention in the last decades, [32, 73, 67, 19, 68, 15] to mention only a few. As far as we know, the existing literature assume either that the state space is a bounded interval instead of the whole  $(0, \infty)$ , as in [32, 5], or that the fragmentation rate has at most a polynomial growth, as in [19, 2, 68, 13]. We can consider here unbounded state space and we relax the latter condition by not assuming any upper bound on the division rate. We obtain thus the existence of the Perron eigentriplet for super-polynomial fragmentation rates. Second, an explicit spectral gap was known only in the case of a constant division rate [73, 59, 20, 10, 68, 84]. Our method allows us to get it for more general fragmentation rates. Finally, it guarantees exponential convergence for measure solutions, thus drastically improving [28].

Let us now state the main result of this section.

**Theorem 5.3.** *Let  $k > 1$  and  $V(x) = 1 + x^k$ . Under the above assumptions, there exists of a unique triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$  of Perron-Frobenius eigenelements with  $\gamma(h) = \|h\|_{\mathcal{B}(V)} = 1$ , i.e. satisfying*

$$\mathcal{L}h = \lambda h \quad \text{and} \quad \gamma(\mathcal{L}f) = \lambda\gamma(f)$$

for all  $f \in C_c^1([0, \infty))$ , where  $\mathcal{L} : C^1([0, \infty)) \rightarrow C^0([0, \infty))$  is defined by

$$\mathcal{L}f(x) = f'(x) + B(x) \left( \int_0^1 f(zx) \wp(dz) - f(x) \right).$$

Moreover there exist constants  $C, \omega > 0$  such that for all  $\mu \in \mathcal{M}(V)$  the solution to Equation (5.5) with  $u_0 = \mu$  satisfies for all  $t \geq 0$ ,

$$\|e^{-\lambda t} u_t - u_0(h)\gamma\|_{\mathcal{M}(V)} \leq C e^{-\omega t} \|u_0\|_{\mathcal{M}(V)}. \quad (5.7)$$

The constants  $\lambda, \omega$ , and  $C$  can be estimated quantitatively, and the eigenfunction  $h$  satisfies  $(1+x)^{1-q(k-1)} \lesssim h \lesssim (1+x)^k$ , for some explicit  $q > 0$ . Note also that we cannot expect the convergence (5.7) to hold true in  $\mathcal{M}(h)$  in general. In particular it is known to be wrong when  $B$  is bounded [12].

The end of the section is devoted to the proof of Theorem 5.3. We can associate to Equation (5.5) a semigroup  $(M_t)_{t \geq 0}$ , to which we will apply our result in Theorem 2.1 after having checked that it verifies Assumption **A** with the functions

$$V(x) = 1 + x^k \quad \text{and} \quad \psi(x) = \frac{1}{2}(1 + x)$$

with  $k > 1$ . The factor  $\frac{1}{2}$  in the definition of  $\psi$  is to satisfy the inequality  $\psi \leq V$ . We only give here the definition of this semigroup as well as its main properties which are useful to verify Assumption **A**, and we refer to Appendix 6.3 for the proofs. For any  $f : [0, \infty) \rightarrow \mathbb{R}$  measurable and locally bounded on  $[0, \infty)$ , we define the family  $(M_t f)_{t \geq 0}$  as the unique solution to the equation

$$M_t f(x) = f(x+t) e^{-\int_0^t B(x+s) ds} + \int_0^t e^{-\int_0^s B(x+s') ds'} B(x+s) \int_0^1 M_{t-s} f(z(x+s)) \wp(dz) ds.$$

This semigroup is positive and preserves  $C^1([0, \infty))$ . More precisely if  $f \in C^1([0, \infty))$ , then the function  $(t, x) \mapsto M_t f(x)$  is continuously differentiable on  $[0, \infty)^2$  and

$$\partial_t M_t f(x) = \mathcal{L} M_t f(x) = M_t \mathcal{L} f(x),$$

where  $\mathcal{L}$  is defined in Theorem 5.3. The space  $\mathcal{B}(V)$  is invariant under  $(M_t)_{t \geq 0}$  and for any  $\mu \in \mathcal{M}(V)$ , we can define by duality  $\mu M_t \in \mathcal{M}(V)$ . The family  $(\mu M_t)_{t \geq 0}$  is solution to (5.5) with initial data  $\mu$ , in a weak sense made precise in Appendix 6.3.

Let  $x_0 \geq 0$  and  $\underline{B} > 0$  such that for all  $x \geq x_0$ ,  $B(x) \geq \underline{B}$ . Now define

$$t_0 = \frac{1 + z_0 + (1 + \epsilon)x_0}{1 - z_0} + \frac{1}{2}, \quad t_1 = \frac{1 - z_0}{2z_0}, \quad \tau = t_0 + t_1, \quad (5.8)$$

and for all integer  $n \geq 0$ ,

$$y_n = \left( \frac{1 + z_0}{2z_0} \right)^n + x_0.$$

**Lemma 5.4.** *i) Setting  $\varphi(x) = 1 - \sqrt{x} + x$ , we have  $\psi \leq \varphi \leq 2\psi$  and there exist  $\zeta > 0$  and  $a < b < \xi$  such that*

$$\mathcal{L}V \leq aV + \zeta\psi, \quad \mathcal{L}\psi \geq b\psi, \quad \mathcal{L}\varphi \leq \xi\varphi,$$

where  $\mathcal{L}$  is the generator of  $(M_t)_{t \geq 0}$  in the sense defined in Section 2.2.

*ii) For all  $n \geq 0$ , all  $x \in [0, y_n]$ , and all  $f : [0, \infty) \rightarrow [0, \infty)$  locally bounded we have*

$$M_\tau f(x) \geq e^{-\tau B(y_n + \tau)} \frac{(c_0 \underline{B})^{n+1} t_1^n}{1 - z_0} \nu(f),$$

where  $\nu$  is the probability measure defined by

$$\nu(f) = \int_{z_0(y_0 + \tau)}^{z_0(y_0 + \tau) + 1} f(y) dy.$$

*iii) For all  $\eta > 0$  there exists  $c_\eta > 0$  such that for all  $t, x \geq 0$  and  $y \in [\eta x, x]$*

$$c_\eta \leq \frac{M_t \psi(y)}{M_t \psi(x)} \leq 1.$$

*iv) For all  $n \geq 0$ , there exists  $d > 0$  such that*

$$d \frac{M_t \psi(x)}{\psi(x)} \leq \frac{M_t \psi(y)}{\psi(y)}$$

for all  $t \geq 0$ ,  $x \in [0, y_n]$  and  $y \in [z_0(y_0 + \tau), z_0(y_0 + \tau) + 1] = \text{supp } \nu$ .

*Proof of Lemma 5.4 i).* Since by Proposition 6.3 the identity  $\partial_t M_t = M_t \mathcal{L}$  is valid for all  $C^1$  functions and the semigroup  $M$  is positive, we only need to prove that  $\mathcal{L}V \leq aV + \zeta\psi$ ,  $\mathcal{L}\psi \geq b\psi$ ,  $\mathcal{L}\varphi \leq \xi\varphi$ . First,

$$\mathcal{L}x^r = rx^{r-1} + (\wp_r - 1)B(x)x^r$$

for any  $r \geq 0$ . We deduce that

$$2\mathcal{L}\psi(x) = 1 + (\wp_0 - 1)B(x) \geq 0,$$

so that  $b = 0$  suits. For  $\varphi$  we have

$$\mathcal{L}\varphi(x) = 1 - \frac{1}{2\sqrt{x}} + (\wp_0 - 1)B(x) - (\wp_{\frac{1}{2}} - 1)B(x)\sqrt{x}.$$

Since  $x \mapsto (\wp_0 - 1) - (\wp_{\frac{1}{2}} - 1)\sqrt{x}$  is negative for  $x > \left(\frac{\wp_0 - 1}{\wp_{\frac{1}{2}} - 1}\right)^2$  and  $B$  is increasing we deduce

$$\mathcal{L}\varphi(x) \leq 1 + (\wp_0 - 1)B\left(\left(\frac{\wp_0 - 1}{\wp_{\frac{1}{2}} - 1}\right)^2\right) =: \frac{\xi}{2} \leq \xi\varphi(x).$$

For  $V$  we have

$$\mathcal{L}V(x) = kx^{k-1} + (\wp_0 - 1)B(x) + (\wp_k - 1)B(x)x^k = \underbrace{\left[ ((\wp_k - 1) + (\wp_0 - 1)x^{-k})B(x) + \frac{k}{x} \right]}_{\rightarrow l := (\wp_k - 1) \lim_{x \rightarrow +\infty} B(x) \text{ when } x \rightarrow +\infty} x^k.$$

Since  $\wp_k < 1$  and  $B$  is increasing, the limit  $l$  belongs to  $[-\infty, 0)$  and we can find  $x_1 > 0$  such that for all  $x \geq x_1$ ,

$$\mathcal{L}V(x) \leq ax^k = aV(x) - a,$$

where  $a = \max\{l/2, -1\} < 0$ . For all  $x \in [0, x_1]$  we have

$$\mathcal{L}V(x) \leq kx_1^{k-1} + (\wp_0 - 1)B(x_1)$$

and finally setting  $\zeta = 2(kx_1^{k-1} + (\wp_0 - 1)B(x_1) - a)$ , we get that for all  $x \geq 0$

$$\mathcal{L}V(x) \leq aV(x) + \frac{\zeta}{2} \leq aV(x) + \zeta\psi(x).$$

It ends the proof of *i*). □

Before proving *ii*), let us briefly comment on the definition of  $t_0, t_1$  and  $y_n$ . The time  $t_1$  and the sequence  $y_n$  are chosen in such a way that

$$y_0 > 0, \quad y_0 \geq x_0, \quad \lim_{n \rightarrow \infty} y_n = +\infty, \quad \text{and} \quad z_0(y_{n+1} + t_1) \leq y_n.$$

The choice of  $t_0$  appears in the proof of the case  $n = 0$  and the definition of  $\nu$ .

Since  $\tau$  is independent of  $n$  and  $y_n \rightarrow +\infty$  when  $n \rightarrow \infty$  we can find  $R$  and  $n$  large enough so that  $\text{supp } \nu \subset K \subset [0, y_n]$ , where  $K = \{x, V(x) \leq R\psi(x)\}$ , and thus *ii*) guarantees that Assumption (A3) is satisfied with time  $\tau$  on  $K$ . More precisely it suffices to take  $R$  and  $n$  large enough so that

$$\frac{1 + (z_0(y_0 + \tau) + 1)^k}{1 + z_0(y_0 + \tau) + 1} \leq \frac{R}{2} \leq \frac{1 + y_n^k}{1 + y_n}. \quad (5.9)$$

*Proof of Lemma 5.4 ii*). Let  $f \geq 0$ . We prove by induction on  $n$  that for all  $x \in [0, y_n]$  and all  $t \in [0, t_1]$  we have

$$M_{t_0+t}f(x) \geq e^{-(t_0+t)B(y_n+\tau)} \frac{(c_0 B)^{n+1}}{1 - z_0} \frac{t^n}{n!} \nu(f), \quad (5.10)$$

which yields the desired result by taking  $t = t_1$ .



We start with the case  $n = 0$ . The Duhamel formula

$$M_t f(x) = f(x+t) e^{-\int_0^t B(x+s) ds} + \int_0^t e^{-\int_0^s B(x+s') ds'} B(x+s) \int_0^1 M_{t-s} f(z(x+s)) \wp(dz) ds$$

ensures, using the positivity of  $M_t$  and the growth of  $B$ , that for all  $t, x \geq 0$

$$M_t f(x) \geq e^{-tB(x+t)} \int_0^t B(x+s) \int_0^1 f(z(x+s) + t-s) \wp(dz) ds.$$

Thus for  $t \geq x_0$  we have for all  $x \geq 0$ , using Assumption (5.6) for the second inequality,

$$\begin{aligned} M_t f(x) &\geq e^{-tB(x+t)} \underline{B} \int_{x_0}^t \int_0^1 f(z(x+s) + t-s) \wp(dz) ds \\ &\geq e^{-tB(x+t)} \underline{B} \frac{c_0}{1-z_0} \int_{z_0(x+t)}^{(z_0-\epsilon)(x+x_0)+t-x_0} f(y) dy. \end{aligned}$$

We deduce that for  $t \in [t_0, t_0 + t_1]$  and  $x \in [0, y_0]$

$$M_t f(x) \geq e^{-tB(x_0+\tau)} \underline{B} \frac{c_0}{1-z_0} \int_{z_0(x_0+t_0+t_1)}^{(z_0-\epsilon)x_0+t_0-x_0} f(y) dy.$$

The time  $t_0$  has been defined in such a way that  $(z_0-\epsilon)x_0+t_0-x_0 = z_0(x_0+t_0+t_1)+1$  so  $\int_{z_0(x_0+t_0+t_1)}^{(z_0-\epsilon)x_0+t_0-x_0} f(y) dy = \nu(f)$  and this finishes the proof of the case  $n = 0$ .

Assume that (5.10) holds for  $n$  and let's check it for  $n+1$ . By Duhamel formula, using that  $y_n \geq x_0$  and  $z_0(y_{n+1} + t_1) \leq y_n$ , we get for  $x \in [y_n, y_{n+1}]$  and  $t \in [0, t_1]$ ,

$$\begin{aligned} M_{t_0+t} f(x) &\geq \int_0^t e^{-\int_0^s B(x+s') ds'} B(x+s) \int_0^1 M_{t_0+t-s} f(z(x+s)) \wp(dz) ds \\ &\geq \underline{B} \int_0^t e^{-sB(x_{n+1}+t_1)} \int_0^{z_0} M_{t_0+t-s} f(z(x+s)) \wp(dz) ds \\ &\geq \underline{B}^{n+2} \frac{c_0^{n+1}}{1-z_0} \nu(f) \int_0^t e^{-sB(x_{n+1}+t_1)} e^{-(t_0+t-s)B(x_n+\tau)} \frac{(t-s)^n}{n!} \int_0^{z_0} \wp(dz) ds \\ &\geq e^{-(t_0+t)B(y_{n+1}+\tau)} \underline{B}^{n+2} \frac{c_0^{n+2}}{1-z_0} \frac{t^{n+1}}{(n+1)!} \nu(f) \end{aligned}$$

and the proof is complete.  $\square$

We now turn to the proof of *iii*) and *iv*), which uses the monotonicity results proved in Lemma 6.5, see Appendix 6.3.

*Proof of Lemma 5.4 iii).* The second inequality readily follows from Lemma 6.5 *ii*). For the first one, we start with a technical result on  $\wp$ . Due to the assumption we made on  $\wp$ , if we set  $z_1 > \max(z_0, 1 - c_0(z_0 - \epsilon/2))$ , we have

$$\varrho := \int_{z_1}^1 \wp(dz) \leq \frac{1}{z_1} \left( 1 - \int_0^{z_1} z \wp(dz) \right) \leq \frac{1 - c_0(z_0 - \epsilon/2)}{z_1} < 1.$$

Using Lemma 6.5 *ii*) and *iii*), we deduce that for all  $t \geq s \geq 0$  and all  $x \geq 0$

$$\int_0^1 M_{t-s}\psi(z(x+s)) \wp(dz) \leq \int_0^1 M_t\psi(zx) \wp(dz) \leq \wp_0 M_t\psi(z_1x) + \varrho M_t\psi(x).$$

Now from the Duhamel formula we get, using that  $t \mapsto t e^{-\int_0^t B(s) ds}$  is bounded on  $[0, \infty)$ ,

$$\begin{aligned} M_t\psi(x) &= \psi(x+t)e^{-\int_0^t B(x+s) ds} + \int_0^t e^{-\int_0^s B(x+s') ds'} B(x+s) \int_0^1 M_{t-s}\psi(z(x+s)) \wp(dz) ds \\ &\leq (1+x+t)e^{-\int_0^t B(s) ds} + \left(1 - e^{-\int_0^t B(x+s) ds}\right) \left(\wp_0 M_t\psi(z_1x) + \varrho M_t\psi(x)\right) \\ &\leq C_0\psi(x) + \wp_0 M_t\psi(z_1x) + \varrho M_t\psi(x). \end{aligned}$$

Choosing an integer  $n$  such that  $z_1^n \leq \eta$  we obtain

$$M_t\psi(x) \leq \frac{C_0}{1-\varrho} \sum_{k=0}^{n-1} \left(\frac{\wp_0}{1-\varrho}\right)^k \psi(x) + \left(\frac{\wp_0}{1-\varrho}\right)^n M_t\psi(\eta x) = C_1\psi(x) + C_2 M_t\psi(\eta x)$$

and since  $M_t\psi(\eta x) \geq \psi(\eta x) \geq \eta\psi(x)$  we obtain for any  $y \in [\eta x, x]$ ,

$$\frac{M_t\psi(y)}{M_t\psi(x)} \geq \frac{M_t\psi(\eta x)}{C_1\psi(x) + C_2 M_t\psi(\eta x)} \geq \frac{\eta}{C_1 + C_2\eta},$$

which ends the proof.  $\square$

*Proof of Lemma 5.4 iv).* Point *iii*) applied with  $\eta = z_0(y_0 + \tau)/y_n$  gives, together with Lemma 6.5 *ii*), that for all  $x \in [0, y_n]$  and  $y \in [z_0(y_0 + \tau), z_0(y_0 + \tau) + 1]$ ,

$$M_t\psi(y) \geq M_t\psi(z_0(y_0 + \tau)) \geq c_\eta M_t\psi(y_n) \geq c_\eta M_t\psi(x).$$

Besides,  $\psi(x)/\psi(y) = (1+x)/(1+y) \geq 1/(z_0(y_0 + \tau) + 2)$  and we get

$$\frac{M_t\psi(y)}{\psi(y)} \geq \frac{c_\eta}{z_0(y_0 + \tau) + 2} \frac{M_t\psi(x)}{\psi(x)},$$

which ends the proof.  $\square$

We are now in position to prove Theorem 5.3.

*Proof of Theorem 5.3.* Fix  $\tau$  defined in (5.8). In Lemma 5.4 *i*) we have verified the assumptions of Proposition 2.2, so we can find a real  $R > 0$  and an integer  $n \geq 0$  large enough so that (5.9) and Assumptions (A1)-(A2) are satisfied with  $K = \{V \leq R\psi\}$ . Then, points *ii*) and *iv*) in Lemma 5.4 ensure that Assumptions (A3) and (A4) are also satisfied. So Assumption **A** is verified for  $(V, \psi)$  and by virtue of Theorem 2.1 inequality (5.7) is proved, as well as the bounds on  $h$ . It remains to check that

$h$  is continuously differentiable and that  $h$  and  $\gamma$  satisfy the eigenvalue equations  $\mathcal{L}h = \lambda h$  and  $\gamma\mathcal{L} = \lambda\gamma$ . By definition of  $h$ , the Duhamel formula gives

$$h(x)e^{\lambda t} = h(x+t)e^{-\int_0^t B(x+s)ds} + \int_0^t e^{-\int_0^s B(x+s')ds'} B(x+s) \int_0^1 e^{\lambda(t-s)} h(z(x+s)) \wp(dz) ds$$

and we deduce that for any  $x > 0$  the function  $t \mapsto h(t+x)$  is continuous and then continuously differentiable. Moreover, we have the identity  $\partial_t M_t h = M_t \mathcal{L}h$  and since  $M_t h = e^{\lambda t} h$  we deduce  $\mathcal{L}h = \lambda h$ .

For the equation on  $\gamma$ , we start from Proposition 6.4 which ensures that

$$e^{\lambda t} \gamma(f) = (\gamma M_t)(f) = \gamma(f) + \int_0^t (\gamma M_s)(\mathcal{L}f) ds = \gamma(f) + \frac{e^{\lambda t} - 1}{\lambda} \gamma(\mathcal{L}f).$$

for any  $f \in C_c^1([0, \infty))$ . Differentiating with respect to  $t$  yields the result.  $\square$

### 5.3 Comments and a few perspectives

First, as for the Harris conservative case [48], our proof relies on a (quantitative) contraction method for the discrete time semigroup  $(M_{n\tau})_{n \geq 0} = (M_\tau^n)_{n \geq 0}$ . It has been extended easily to the continuous setting. We can thus actually state analogous results for a discrete time semigroup  $(M^n)_{n \in \mathbb{N}}$  by making the following assumption for a couple of positive functions  $(V, \psi)$  with  $\psi \leq V$ .

**Assumption B.** There exist some integer  $\tau > 0$ , real numbers  $\beta > \alpha > 0$ ,  $\theta > 0$ ,  $(c, d) \in (0, 1]^2$ , some set  $K \subset \mathcal{X}$  such that  $\sup_K V/\psi < \infty$ , and some probability measure  $\nu$  on  $\mathcal{X}$  supported by  $K$  such that

$$(B1) \quad M^\tau V \leq \alpha V + \theta \mathbf{1}_K \psi,$$

$$(B2) \quad M^\tau \psi \geq \beta \psi,$$

$$(B3) \quad \inf_{x \in K} \frac{M^\tau(f\psi)(x)}{M^\tau \psi(x)} \geq c \nu(f) \quad \text{for all } f \in \mathcal{B}_+(V/\psi),$$

$$(B4) \quad \nu \left( \frac{M^{n\tau} \psi}{\psi} \right) \geq d \sup_{x \in K} \frac{M^{n\tau} \psi(x)}{\psi(x)} \quad \text{for all positive integers } n.$$

The counterpart of Theorem 2.1 is stated below.

**Theorem 5.5.** *i) Let  $(V, \psi)$  be a couple of measurable functions from  $\mathcal{X}$  to  $(0, \infty)$  which satisfies Assumption B. Then, there exists a unique triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$  of eigenelements of  $M$  with  $\gamma(h) = \|h\|_{\mathcal{B}(V)} = 1$ , i.e. satisfying*

$$\gamma M = \lambda \gamma \quad \text{and} \quad M h = \lambda h. \quad (5.11)$$

Moreover, there exists  $C > 0$  and  $\rho \in (0, 1)$  such that for all  $n \geq 0$  and  $\mu \in \mathcal{M}(V)$ ,

$$\|\lambda^{-n} \mu M^n - \mu(h)\gamma\|_{\mathcal{M}(V)} \leq C \|\mu\|_{\mathcal{M}(V)} \rho^{-n}. \quad (5.12)$$

ii) Assume that there exist a positive measurable function  $V$ , a triplet  $(\gamma, h, \lambda) \in \mathcal{M}_+(V) \times \mathcal{B}_+(V) \times \mathbb{R}$ , and constants  $C, \rho > 0$  such that (5.11) and (5.12) hold. Then, the couple  $(V, h)$  satisfies Assumption **B**.

In addition, we recover the various bounds on the eigenvector as in Lemma 3.4 or on the eigenvalue as in (3.8). We also recover that if  $\inf_{\mathcal{X}} V > 0$  then there exists  $C > 0$  and  $\pi \in \mathcal{P}(V)$  such that for all  $\mu \in \mathcal{P}(V)$  and  $n \geq 0$ ,

$$\left\| \frac{\mu M^n}{\mu M^n \mathbf{1}} - \pi \right\|_{\text{TV}} \leq C \frac{\mu(V)}{\mu(h)} \rho^n.$$

As a consequence, Assumption **B** gives sufficient conditions to have the existence, uniqueness and convergence to a QSD for a Markov chain. Moreover, the convergence of the  $Q$ -process, the description of the domain of attraction and the bounds on the extinction times can be then obtained by usual procedure, see *e.g.* [22, 80].

Second, for the sake of simplicity, we have not allowed  $\psi$  to vanish in this paper. This excludes reducible structures. To illustrate this fact, let us consider the case where  $\mathcal{X} = \{1, 2\}$  similar to [11, Example 3.5] where

$$M = \begin{pmatrix} a & b \\ 0 & c \end{pmatrix},$$

for  $a, b, c > 0$ . In any case, Theorem 5.5 for  $\mathcal{X} = \{2\}$  implies that if  $\mu(\{1\}) = 0$  then  $c^{-n} \mu M^n$  tends to  $\delta_2$ . Here it is an equality, but it can be easily generalized into a reducible example on two subspaces. Now if  $\mu(\{1\}) > 0$  then one can use Theorem 5.5 only when  $c > a$ . Indeed, one can choose  $\psi$  to be the right eigenvector:

$$\psi = (b, c - a) = b \mathbf{1}_{\{1\}} + (c - a) \mathbf{1}_{\{2\}}$$

and show in this case, that, up to a renormalisation,  $\mu M^n$  converges to  $\delta_2$ . When  $c \leq a$ , one can not use Theorem 5.5 and indeed there is no positive right eigenvector. However, allowing  $\psi$  to vanish enables to treat the case  $c < a$ , as in [22, Section 6]. Focusing on initial measures  $\mu$  such that  $\mu(\psi) > 0$ , a large part of our results actually holds when  $\psi \geq 0$ . Indeed (B2) gives that if  $\mu(\psi) > 0$  then  $\mu M^n \psi > 0$  for every  $n \geq 0$ .

Up to our knowledge, the critical case  $a = c$  remains a challenging issue. In this case, there is no spectral gap. Nevertheless, we believe that our approach could be extended to this case by allowing  $n$ -dependent constant  $d = d_n$  in (B4) to vary with time as in [11, Assumption (**H**<sub>4</sub>)].

Finally, we recall that the results obtained here rely on a contraction method. Several extensions to the non-homogeneous setting are expected, in the same vein as [8] for time inhomogeneous linear PDEs. One can now relax the “coming down from infinity” property imposed by the generalized Doeblin condition of [8]. Similarly, let us recall that the expectation of a branching process yields the first moment semigroup, which usually drives the extinction of the process (criticality) and provides its deterministic renormalization (Kesten-Stigum theorem). The method of this paper should provide a powerful tool to analyse the first moment semigroup of a branching process with infinite number of types, including in varying environment, see [6, 7, 61] for some motivations in population dynamics and queuing systems. We also mention that time inhomogeneity provides a natural point of view to deal with non-linearity in large population approximations of systems with interaction. These points should be partially addressed in forthcoming works.

## 6 Appendix

### 6.1 Conservative operators

The following useful result is a direct generalization of the contraction result of Hairer and Mattingly [48], in the same vein as [61, 22]. We consider a positive operator  $P$  acting both on bounded measurable functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  on the right and on bounded measures  $\mu$  of finite mass on the left, and such that  $(\mu P)f = \mu(Pf)$ . Note that the right action of  $P$  extends trivially to any measurable function  $f : \mathcal{X} \rightarrow [0, +\infty]$ . We assume that  $P$  is conservative in the sense that  $P\mathbf{1} = \mathbf{1}$ . In other words, if  $\mu$  is a probability measure, then so does  $\mu P$ .

**Theorem 6.1.** *Assume that there exist two measurable functions  $\mathcal{V}, \mathcal{W} : \mathcal{X} \rightarrow [0, \infty)$ ,  $(\mathbf{a}, \mathbf{b}) \in (0, 1)^2$ ,  $\mathbf{c} > 0$ ,  $\mathfrak{R} > 2\mathbf{c}/(1 - \mathbf{a})$  and a probability measure  $\nu$  on  $\mathcal{X}$  such that:*

- for all  $x \in \mathcal{X}$ ,

$$P\mathcal{V}(x) \leq \mathbf{a}\mathcal{W} + \mathbf{c}, \tag{6.1}$$

- for all  $x \in \{\mathcal{W} \leq \mathfrak{R}\}$ ,

$$\delta_x P \geq \mathbf{b}\nu. \tag{6.2}$$

Then, there exist  $\mathfrak{h} \in (0, 1)$  and  $\kappa > 0$  such that for all probability measures  $\mu_1, \mu_2$ ,

$$\|\mu_1 P - \mu_2 P\|_{\mathcal{M}(1+\kappa\mathcal{V})} \leq \mathfrak{h} \|\mu_1 - \mu_2\|_{\mathcal{M}(1+\kappa\mathcal{W})}.$$

In particular, for any  $\mathbf{b}' \in (0, \mathbf{b})$  and  $\mathbf{a}' \in (\mathbf{a} + 2\mathbf{c}/\mathfrak{R}, 1)$ , one can choose

$$\kappa = \frac{\mathbf{b}'}{\mathbf{c}}, \quad \eta = \max \left\{ 1 - (\mathbf{b} - \mathbf{b}'), \frac{2 + \kappa\mathfrak{R}\mathbf{a}'}{2 + \kappa\mathfrak{R}} \right\}.$$

Usually, for conservative semigroups and Markov chains [65, 48], Theorem 6.1 is stated and used with one single function  $\mathcal{V} = \mathcal{W}$ . For the sake of completeness, we briefly give the proof of Theorem 6.1, which is an adaptation of [48, Theorem 1.3]. For any function  $V : \mathcal{X} \rightarrow [0, \infty)$ , let us define the following distance on  $\mathcal{X}$ :

$$\text{dist}_V(x, y) = \begin{cases} 0 & x = y \\ 2 + V(x) + V(y) & x \neq y \end{cases}$$

We also introduce a semi-norm on measurable functions  $f : \mathcal{X} \rightarrow \mathbb{R}$  defined by

$$\|f\|_{\text{Lip}(V)} = \sup_{x \neq y} \frac{|f(x) - f(y)|}{\text{dist}_V(x, y)},$$

and the associated (Wasserstein) metric on the set of probability measures given by

$$\text{dist}_V(\mu_1, \mu_2) = \sup_{\|f\|_{\text{Lip}(V)} \leq 1} \int_{\mathcal{X}} f(x)(\mu_1 - \mu_2)(dx) = \|\mu_1 - \mu_2\|_{\mathcal{M}(1+V)},$$

using [48, Lemma 2.1]. We prove now Theorem 6.1, which is a simple adaptation of the proofs of Theorems 1.3 and 3.1 in [48].

*Proof of Theorem 6.1.* Let  $f$  be a test function such that  $\|f\|_{\text{Lip}(\kappa\mathcal{V})} \leq 1$  and take  $x \neq y$ . Fix  $\mathbf{a}' \in (\mathbf{a} + 2\mathbf{c}/\mathfrak{R}, 1)$  and  $\mathbf{b}' \in (0, \mathbf{b})$ , and set  $\kappa = \mathbf{b}'/\mathbf{c}$  and

$$\eta = \max \{ 1 - \mathbf{b} + \mathbf{b}', (2 + \kappa\mathfrak{R}\mathbf{a}')/(2 + \kappa\mathfrak{R}) \}.$$

Considering successively the cases  $\mathcal{W}(x) + \mathcal{W}(y) \geq \mathfrak{R}$  and  $\mathcal{W}(x) + \mathcal{W}(y) \leq \mathfrak{R}$  as in [48, Proof of Theorem 3.1], and using (6.1) and (6.2), we obtain

$$|Pf(x) - Pf(y)| \leq \eta \text{dist}_{\kappa\mathcal{W}}(x, y),$$

so that  $\|Pf\|_{\text{Lip}(\kappa\mathcal{W})} \leq \eta$ . Finally, for any probability measure  $\mu_1$  and  $\mu_2$ ,

$$\begin{aligned} \text{dist}_{\kappa\mathcal{V}}(\mu_1 P, \mu_2 P) &= \sup_{\|f\|_{\text{Lip}(\kappa\mathcal{V})} \leq 1} \int_{\mathcal{X}} Pf(x)(\mu_1 - \mu_2)(dx) \\ &\leq \sup_{\|Pf\|_{\text{Lip}(\kappa\mathcal{W})} \leq \eta} \int_{\mathcal{X}} Pf(x)(\mu_1 - \mu_2)(dx) = \eta \text{dist}_{\kappa\mathcal{W}}(\mu_1, \mu_2) \end{aligned}$$

and the proof is complete.  $\square$

## 6.2 Localization argument

We detail here how the drift conditions (5.1)-(5.2) ensure (5.3)-(5.4), namely

$$\begin{aligned} \mathbb{E}_n[V(X_t)] &\leq V(x) + \int_0^t \mathbb{E}_n[(aV + \zeta\psi)(X_s)]ds, \\ \psi(x) + \int_0^t \mathbb{E}_n[b\psi(X_s)]ds &\leq \mathbb{E}_x[\psi(X_t)] \leq \psi(x) + \int_0^t \mathbb{E}_n[\xi\psi(X_s)]ds, \end{aligned}$$

for all  $n \geq 1$  and  $t \geq 0$ . Following [66], for  $m \geq 1$ , we let  $T_m = \inf\{t > 0 : X_t \geq m\}$  and  $(X_t^m)_{t \geq 0}$  be the Markov process defined by  $X_t^m = X_t \mathbf{1}_{t < T_m}$ . We extend functions  $V, \psi$  on  $\mathbb{N}$  by setting  $V(0) = \psi(0) = 0$ . Using (5.1), (5.2),  $V(0) \leq V(m)$  and writing  $O_m = \{0, 1, \dots, m-1\}$ , the strong generator  $\mathcal{L}^m$  of  $X^m$  satisfies

$$\begin{aligned} \mathcal{L}^m V &\leq aV + \zeta\psi \quad \text{and} \quad \mathcal{L}^m \psi \leq \xi\psi \quad \text{on } O_m \\ \mathcal{L}^m \psi(m-1) &= \mathcal{L}\psi(m-1) - b\psi(m) \geq b(1-\eta)\psi(m-1) \\ \mathcal{L}^m \psi &\geq b\psi \quad \text{on } O_{m-1}. \end{aligned}$$

First, using  $\mathcal{L}^m V \leq (a + \zeta)V$  on  $O_m$  and  $V(n) \rightarrow \infty$  as  $n \rightarrow \infty$ , [66, Theorem 2.1] ensures that  $\lim_{m \rightarrow \infty} T_m = \infty$  and

$$\mathbb{E}_n[V(X_t)] \leq e^{(a+\zeta)t}V(n)$$

for every  $n \in \mathbb{N}$ . Second  $\mathcal{L}^m \psi \leq \xi\psi$  on  $O_m$  and  $\psi$  is bounded on  $O_m$ . Using that  $X^m$  coincides with  $X$  on  $[0, T_m)$ , Fatou's lemma and Kolmogorov equation give

$$\begin{aligned} \mathbb{E}_n[\psi(X_t)] &\leq \liminf_{m \rightarrow \infty} \mathbb{E}_x[\psi(X_t^m)] \\ &= \psi(n) + \liminf_{m \rightarrow \infty} \mathbb{E}_n \left[ \int_0^{t \wedge T_m} \mathcal{L}^m \psi(X_s) ds \right] \leq \psi(x) + \xi \int_0^t \mathbb{E}_n[\psi(X_s)] ds. \end{aligned}$$

Moreover  $\psi(X_t^m) = \mathbf{1}_{t \leq T_m} \psi(X_t) \leq \psi(X_t)$  and  $X_t^m \rightarrow X_t$  as  $m \rightarrow \infty$ . Using  $\mathcal{L}^m \psi \geq b\psi - b\eta\psi \mathbf{1}_{m-1}$  on  $O_m$  and bounded convergence twice yields  $\mathbb{E}(\int_0^t \psi(X_s) \mathbf{1}_{X_s=m-1} ds) \rightarrow 0$  as  $m \rightarrow \infty$  and

$$\begin{aligned} \mathbb{E}_n[\psi(X_t)] &= \lim_{m \rightarrow \infty} \mathbb{E}_x[\psi(X_t^m)] \\ &= \psi(n) + \lim_{m \rightarrow \infty} \mathbb{E}_n \left[ \int_0^{t \wedge T_m} \mathcal{L}^m \psi(X_s) ds \right] \geq \psi(x) + b \int_0^t \mathbb{E}_n[\psi(X_s)] ds. \end{aligned}$$

Using Fatou's lemma as above for  $V$  ends the proof of (5.3)-(5.4).

## 6.3 The growth-fragmentation semigroup

We give here the details of the construction of the growth-fragmentation semigroup and prove its basic properties, along the lines of [38, 44, 45]. For a function  $f \in$

$\mathcal{B}_{loc}([0, \infty))$ , we define the family  $(M_t f)_{t \geq 0} \subset \mathcal{B}_{loc}([0, \infty))$  through the Duhamel formula

$$M_t f(x) = f(x+t)e^{-\int_0^t B(x+s)ds} + \int_0^t e^{-\int_0^s B(x+s')ds'} B(x+s) \int_0^1 M_{t-s} f(z(x+s)) \wp(dz) ds.$$

We first prove that this indeed defines uniquely the family  $(M_t f)_{t \geq 0}$ . Then, we verify that the associated family  $(M_t)_{t \geq 0}$  is a semigroup of linear operators, which provides the unique solution to the growth-fragmentation (5.5) on the space  $\mathcal{M}(V)$  with  $V(x) = 1 + x^k$ ,  $k > 1$ . Finally we provide some useful monotonicity properties for this semigroup, which are consequences of the monotonicity assumption on  $B$ .

**Lemma 6.2.** *For any  $f \in \mathcal{B}_{loc}([0, \infty))$  there exists a unique  $\bar{f} \in \mathcal{B}_{loc}([0, \infty)^2)$  such that for all  $t \geq 0$  and  $x \geq 0$*

$$\bar{f}(t, x) = f(x+t)e^{-\int_0^t B(x+s)ds} + \int_0^t e^{-\int_0^s B(x+s')ds'} B(x+s) \int_0^1 \bar{f}(t-s, z(x+s)) \wp(dz) ds.$$

Moreover if  $f$  is nonnegative/continuous/continuously differentiable, then so does  $\bar{f}$ . In the latter case  $\bar{f}$  satisfies the partial differential equation

$$\partial_t \bar{f}(t, x) = \mathcal{L} \bar{f}(t, x) = \partial_x \bar{f}(t, x) + B(x) \left[ \int_0^1 \bar{f}(t, zx) \wp(dz) - \bar{f}(t, x) \right].$$

*Proof.* Let  $f \in \mathcal{B}_{loc}([0, \infty))$  and define on  $\mathcal{B}_{loc}([0, \infty)^2)$  the mapping  $\Gamma$  by

$$\Gamma g(t, x) = f(x+t)e^{-\int_0^t B(x+s)ds} + \int_0^t e^{-\int_0^s B(x+s')ds'} B(x+s) \int_0^1 g(t-s, z(x+s)) \wp(dz) ds.$$

Now for  $T, A > 0$  define the set  $\Omega_{T,A} = \{(t, x) \in [0, T] \times [0, \infty), x+t < A\}$  and denote by  $\mathcal{B}_b(\Omega_{T,A})$  the Banach space of bounded measurable functions on  $\Omega_{T,A}$ , endowed with the supremum norm  $\|\cdot\|_\infty$ . Clearly  $\Gamma$  induces a mapping  $\mathcal{B}_b(\Omega_{T,A}) \rightarrow \mathcal{B}_b(\Omega_{T,A})$ , still denoted by  $\Gamma$ . To build a fixed point of  $\Gamma$  in  $\mathcal{B}_{loc}([0, \infty)^2)$  we prove that it admits a unique fixed point in any  $\mathcal{B}_b(\Omega_{A,A})$ .

Let  $A > 0$  and  $T < 1/(\wp_0 B(A))$ . For any  $g_1, g_2 \in \mathcal{B}_b(\Omega_{T,A})$  we have

$$\|\Gamma g_1 - \Gamma g_2\|_\infty \leq \wp_0 T B(A) \|g_1 - g_2\|_\infty$$

and  $\Gamma$  is a contraction. The Banach fixed point theorem then guarantees the existence of a unique fixed point  $g_{T,A}$  of  $\Gamma$  in  $\mathcal{B}_b(\Omega_{T,A})$ . The same argument on  $\Omega_{T,A-T}$  ensures that  $g_{T,A}$  can be extended into a unique fixed point  $g_{2T,A}$  of  $\Gamma$  on  $\Omega_{2T,A}$ . Iterating the procedure we finally get a unique fixed point  $g_A$  of  $\Gamma$  in  $\mathcal{B}_b(\Omega_{A,A})$ .

For  $A' > A > 0$  we have  $g_{A'}|_{\Omega_A} = g_A$  by uniqueness of the fixed point in  $\mathcal{B}_b(\Omega_{A'})$ , and we can define  $\bar{f}$  by setting  $\bar{f}|_{\Omega_A} = g_A$  for any  $A > 0$ . Clearly the function  $\bar{f}$



thus defined is the unique fixed point of  $\Gamma$  in  $\mathcal{B}_{loc}([0, \infty)^2)$ . Since  $\Gamma$  preserves the closed cone of nonnegative functions if  $f$  is nonnegative, the fixed point  $\bar{f}$  necessarily belongs to this cone when  $f$  is nonnegative. Similarly, the space  $C([0, \infty)^2)$  of continuous functions being a closed subspace of  $\mathcal{B}_{loc}([0, \infty)^2)$ , the fixed point  $\bar{f}$  is continuous when  $f$  is so.

Consider now that  $f$  is continuously differentiable on  $[0, \infty)$ . The space  $C^1([0, \infty)^2)$  is not closed in  $\mathcal{B}_{loc}([0, \infty)^2)$  for the norm  $\|\cdot\|_\infty$ . For proving the continuous differentiability of  $\bar{f}$  we repeat the fixed point argument in  $\{g \in C^1(\Omega_{T,A}), g(0, \cdot) = f\}$ , endowed with the norm  $\|g\|_{C^1} = \|g\|_\infty + \|\partial_t g\|_\infty + \|\partial_x g\|_\infty$ . Differentiating  $\Gamma g$  with respect to  $t$  we get

$$\partial_t \Gamma g(t, x) = \mathcal{L}f(x+t)e^{-\int_0^t B(x+s)ds} + \int_0^t e^{-\int_0^s B(x+s')ds'} B(x+s) \int_0^1 \partial_t g(t-s, z(x+s)) \wp(dz) ds$$

and differentiating the alternative formulation

$$\Gamma g(t, x) = f(x+t)e^{-\int_x^{x+t} B(y)dy} + \int_x^{x+t} e^{-\int_x^y B(y')dy'} B(y) \int_0^1 g(t+x-y, zy) \wp(dz) dy$$

with respect to  $x$  we obtain

$$\begin{aligned} \partial_x \Gamma g(t, x) &= \mathcal{L}f(x+t)e^{-\int_x^{x+t} B(y)dy} + B(x) \left( f(x+t)e^{-\int_x^{x+t} B(y)dy} - \int_0^1 g(t, zx) \wp(dz) \right) \\ &\quad + B(x) \int_x^{x+t} e^{-\int_x^y B(y')dy'} B(y) \int_0^1 g(t+x-y, zy) \wp(dz) dy \\ &\quad + \int_x^{x+t} e^{-\int_x^y B(y')dy'} B(y) \int_0^1 \partial_t g(t+x-y, zy) \wp(dz) dy \\ &= \left[ \mathcal{L}f(x+t) + B(x)f(x+t) - B(x) \int_0^1 f(zx) \wp(dz) \right] e^{-\int_x^{x+t} B(y)dy} \\ &\quad + \int_x^{x+t} e^{-\int_x^y B(y')dy'} (B(y) - B(x)) \int_0^1 \partial_t g(t+x-y, zy) \wp(dz) dy \\ &\quad + B(x) \int_x^{x+t} e^{-\int_x^y B(y')dy'} \int_0^1 \partial_x g(t+x-y, zy) z \wp(dz) dy. \end{aligned}$$

On the one hand using the second expression of  $\partial_x \Gamma g(t, x)$  above we deduce that for  $g_1, g_2 \in C^1(\Omega_{T,A})$  such that  $g_1(0, \cdot) = g_2(0, \cdot) = f$  we have

$$\|\Gamma g_1 - \Gamma g_2\|_{C^1} \leq 2\wp_0 T B(A) \|g_1 - g_2\|_{C^1}.$$

Thus  $\Gamma$  is a contraction for  $T < 1/(2\wp_0 T B(A))$  and this guarantees that the fixed point  $\bar{f}$  necessarily belongs to  $C^1([0, \infty)^2)$ . On the other hand using the first expression of  $\partial_x \Gamma g(t, x)$  we have

$$\partial_t \Gamma g(t, x) - \partial_x \Gamma g(t, x) = B(x) \left[ \int_0^1 g(t, zx) \wp(dz) - \Gamma g(t, x) \right]$$

and accordingly the fixed point satisfies  $\partial_t \bar{f} = \mathcal{L}\bar{f}$ .  $\square$

With Lemma 6.2 at hand we can define for any  $t \geq 0$  the mapping  $M_t$  on  $\mathcal{B}_{loc}(0, \infty)$  by setting

$$M_t f(x) = \bar{f}(t, x).$$

**Proposition 6.3.** *The family  $(M_t)_{t \geq 0}$  defined above is a positive semigroup of linear operators on  $\mathcal{B}_{loc}([0, \infty))$ . If  $f \in C^1([0, \infty))$  then the function  $(t, x) \mapsto M_t f(x)$  is continuously differentiable and satisfies*

$$\partial_t M_t f(x) = \mathcal{L} M_t f(x) = M_t \mathcal{L} f(x).$$

Additionally for any  $k > 1$  the space  $\mathcal{B}(V)$  with  $V(x) = 1 + x^k$  is invariant under  $(M_t)_{t \geq 0}$ , and for all  $t \geq 0$  the restriction of  $M_t$  to  $\mathcal{B}(V)$  is a bounded operator.

*Proof.* The linearity and the semigroup property readily follow from the uniqueness of the fixed point in Lemma 6.2. The positivity and the stability of  $C^1([0, \infty))$  are direct consequences of Lemma 6.2, as well as the relation  $\partial_t M_t f = \mathcal{L} M_t f$ . For getting the second one  $\partial_t M_t f = M_t \mathcal{L} f$ , it suffices to remark from the computation of  $\partial_t \Gamma g$  in the proof of Lemma 6.2 that  $\partial_t M_t f$  is the unique fixed point of  $\Gamma$  with initial data  $\mathcal{L} f$ . For the invariance of  $\mathcal{B}(V)$  we compute

$$\mathcal{L} V(x) = 1 + kx^{k-1} + (\wp_0 - 1)B(x) + (\wp_k - 1)B(x)x^k$$

which is bounded on  $[0, \infty)$  since  $(\wp_0 - 1)B(x) + (\wp_k - 1)B(x)x^k \leq 0$  when  $x \geq \left(\frac{\wp_0 - 1}{1 - \wp_k}\right)^{\frac{1}{k}}$ . We deduce that there exists  $C > 0$  such that  $\mathcal{L} V \leq CV$  and since  $V \in C^1([0, \infty))$  we get

$$M_t V(x) = V(x) + \int_0^t M_s(\mathcal{L} V)(x) ds \leq e^{Ct} V(x).$$

Positivity of  $M_t$  then yields  $\|M_t f\|_{\mathcal{B}(V)} \leq e^{Ct} \|f\|_{\mathcal{B}(V)}$ .  $\square$

Now we define, for  $t \geq 0$  and  $\mu \in \mathcal{M}_+(V)$ , the positive measure  $\mu M_t$  by setting for any measurable set  $A \subset [0, \infty)$

$$(\mu M_t)(A) = \mu(M_t \mathbf{1}_A).$$

Then, for  $\mu \in \mathcal{M}(V)$  we define  $\mu M_t \in \mathcal{M}(V)$  as the equivalence class of  $(\mu_+ M_t, \mu_- M_t)$ .

**Proposition 6.4.** *The family  $(M_t)_{t \geq 0}$  defined above is a positive semigroup of bounded linear operators on  $\mathcal{M}(V)$ . Moreover for any  $\mu \in \mathcal{M}(V)$  the family  $(\mu M_t)_{t \geq 0}$  is solution to Equation (5.5) in the sense that for all  $f \in C_c^1([0, \infty))$  and all  $t \geq 0$*

$$(\mu M_t)(f) = \mu(f) + \int_0^t (\mu M_s)(\mathcal{L} f) ds.$$

*Proof.* Let  $\mu \in \mathcal{M}(V)$  and  $f \in C_c^1([0, \infty))$ . From Proposition 6.3 we know that  $\partial_t M_t f = M_t \mathcal{L} f$  which gives by integration in time

$$M_t f(x) = f(x) + \int_0^t M_s \mathcal{L} f(x) ds = f(x) + \int_0^t M_s (f' - Bf)(x) ds + \int_0^t M_s \mathcal{F} f(x) ds$$

for all  $x \geq 0$ , where we have set

$$\mathcal{F} f(x) = B(x) \int_0^1 f(zx) \varphi(dz).$$

Since  $f' - Bf \in \mathcal{B}(V)$  we have  $|M_s(f' - Bf)| \leq \|f' - Bf\|_{\mathcal{B}(V)} e^{C_s V}$  and Fubini's theorem ensures that

$$\mu \left( \int_0^t M_s (f' - Bf) ds \right) = \int_0^t (\mu M_s) (f' - Bf) ds.$$

The last term deserves a bit more attention since  $\mathcal{F} f$  can be not bounded by  $V$ . Consider  $g \in C_c^1([0, \infty))$  such that  $g \geq |f|$ . By positivity of  $M_s$  and  $\mathcal{F}$  we have  $|M_s \mathcal{F} f| \leq M_s \mathcal{F} |f| \leq M_s \mathcal{F} g$  and since  $g \in C_c^1(0, \infty)$

$$\mu_{\pm} \left( \int_0^t M_s \mathcal{F} g ds \right) = \mu_{\pm} \left( M_t g - g - \int_0^t M_s (g' - Bg) ds \right) < +\infty.$$

Then  $(s, x) \mapsto M_s \mathcal{F} f(x)$  is  $(ds \times \mu)$ -integrable and Fubini's theorem yields

$$\mu \left( \int_0^t M_s \mathcal{F} f ds \right) = \int_0^t (\mu M_s) (\mathcal{F} f) ds,$$

which ends the proof.  $\square$

We end this appendix by giving some monotonicity results on  $(M_t)_{t \geq 0}$ , which are useful for verifying (A4) in Section 5.2. They are valid under the monotonicity assumption we made on the fragmentation rate  $B$ .

**Lemma 6.5.** *i) For any  $x \geq 0$ ,  $t \mapsto M_t \psi(x)$  is increasing.*

*ii) For any  $t \geq 0$ ,  $x \mapsto M_t \psi(x)$  is increasing.*

*iii) For any  $T > 0$ ,  $z \in [0, 1]$  and  $x \geq 0$ ,  $t \mapsto M_t \psi(z(x + T - t))$  increases on  $[0, T]$ .*

*Proof.* The point *i)* readily follows from  $\partial_t M_t \psi = M_t (\mathcal{L} \psi)$ , since  $M_t$  is positive and  $2\mathcal{L} \psi(x) = 1 + (\varphi_0 - 1)B(x) \geq 0$ .

Let us prove *ii)*. Define  $f(t, x) = \partial_x M_t \psi(x)$  which satisfies

$$\partial_t f(t, x) = \partial_x f(t, x) - B(x) f(t, x) + B(x) \int_0^1 f(t, zx) z \varphi(dz) + C(x),$$

with  $C(x) = -B'(x)M_t\psi(x) + B'(x) \int_0^1 M_t\psi(zx) \varphi(dz)$ . Since  $\partial_t M_t\psi(x) = \mathcal{L}M_t\psi(x)$ ,  $\partial_t M_t\psi(x) \geq 0$ , and  $B' \geq 0$ , we have

$$C(x) = \frac{B'(x)}{B(x)} (\partial_t M_t\psi(x) - \partial_x M_t\psi(x)) \geq -\frac{B'(x)}{B(x)} f(t, x)$$

and as a consequence

$$\partial_t f(t, x) \geq \mathcal{A}f(t, x) := \partial_x f(t, x) - \left( B(x) + \frac{B'(x)}{B(x)} \right) f(t, x) + B(x) \int_0^1 f(t, zx) z \varphi(dz).$$

Similarly to  $\mathcal{L}$  the operator  $\mathcal{A}$  generates a positive semigroup  $(U_t)_{t \geq 0}$ . It is a standard result that it enjoys the following maximum principle  $\partial_t f(t, x) \geq \mathcal{A}f(t, x) \implies f(t, x) \geq U_t f_0(x)$ , where  $f_0 = f(0, \cdot)$ . Since  $f(0, x) = \psi'(x) = \frac{1}{2} \geq 0$  we deduce from the positivity of  $U_t$  that  $f(t, x) \geq 0$  for all  $t, x > 0$ , which ends the proof of *ii*).

We turn to the proof of *iii*). The case  $z = 0$  corresponds to *ii*) and we consider now  $z \in (0, 1]$ . Setting  $f(t, x) = M_t\psi(z(x + T - t))$  we have using *ii*),  $\partial_x f(t, x) = z \partial_x M_t\psi(z(x + T - t)) \geq 0$  and

$$\begin{aligned} \partial_t f(t, x) &= (\partial_t M_t\psi)(z(x + T - t)) - z (\partial_x M_t\psi)(z(x + T - t)) \\ &= \frac{1-z}{z} \partial_x f(t, x) - B(z(x + T - t)) f(t, x) \\ &\quad + B(z(x + T - t)) \int_0^1 f(t, z'x - (1-z')(T-t)) \varphi(dz'). \end{aligned} \quad (6.3)$$

Now define  $g(t, x) = \partial_t f(t, x)$  and differentiate the above equation with respect to  $t$  and use again (6.3) to get

$$\begin{aligned} \partial_t g(t, x) &= \frac{1-z}{z} \partial_x g(t, x) - B(z(x + T - t)) g(t, x) \\ &\quad + B(z(x + T - t)) \int_0^1 g(t, z'x - (1-z')(T-t)) \varphi(dz') \\ &\quad + z \frac{B'}{B}(z(x + T - t)) \left( \frac{1-z}{z} \partial_x f(t, x) - g(t, x) \right) \\ &\quad + B(z(x + T - t)) \int_0^1 (1-z') \partial_x f(t, z'x - (1-z')(T-t)) \varphi(dz') \end{aligned}$$

and using the positivity of  $\partial_x f$ ,  $B$  and  $B'$  we finally obtain

$$\begin{aligned} \partial_t f(t, x) &\geq \frac{1-z}{z} \partial_x g(t, x) - \left( B + z \frac{B'}{B} \right) (z(x + T - t)) g(t, x) \\ &\quad + B(z(x + T - t)) \int_0^1 g(t, z'x - (1-z')(T-t)) \varphi(dz'). \end{aligned}$$

Since  $g(0, x) = \frac{1-z}{2} + \frac{\varphi_0 - 1}{2} B(z(x + T)) \geq 0$  we deduce from the maximum principle that  $g(t, x) \geq 0$ .  $\square$

## Acknowledgments

The authors are grateful to the anonymous referees and the AE for their suggestions and remarks which have improved the previous version. V.B., B.C., and A.M. have received the support of the Chair “Modélisation Mathématique et Biodiversité” of VEOLIA-Ecole Polytechnique-MnHn-FX. The authors have been also supported by ANR projects, funded by the French Ministry of Research: B.C. by ANR MESA (ANR-18-CE40-006), V.B. by ANR ABIM (ANR-16-CE40-0001) and ANR CADENCE (ANR-16-CE32-0007), and A.M. by ANR MEMIP (ANR-16-CE33-0018).

## References

- [1] A. Asselah and M.-N. Thai. A note on the rightmost particle in a Fleming-Viot process. Preprint, arXiv:1212.4168, 2012.
- [2] D. Balagué, J. A. Cañizo, and P. Gabriel. Fine asymptotics of profiles and relaxation to equilibrium for growth-fragmentation equations with variable drift rates. *Kinet. Relat. Models*, 6(2):219–243, 2013.
- [3] J. Banasiak and L. Arlotti. *Perturbations of Positive Semigroups with Applications*. Springer Monographs in Mathematics. Springer-Verlag, London, 2006.
- [4] J. Banasiak, W. Lamb, and P. Laurençot. *Analytic Methods for Coagulation-Fragmentation Models, Volume I*. Chapman & Hall/CRC Monographs and Research Notes in Mathematics. 2019.
- [5] J. Banasiak, K. Pichór, and R. Rudnicki. Asynchronous exponential growth of a general structured population model. *Acta Appl. Math.*, 119:149–166, 2012.
- [6] V. Bansaye. Ancestral lineage and limit theorems for branching Markov chains. *J. Theor. Probab.*, 2018.
- [7] V. Bansaye and A. Camanes. Queueing for an infinite bus line and aging branching process. *Queueing Syst.*, 2018.
- [8] V. Bansaye, B. Cloez, and P. Gabriel. Ergodic Behavior of Non-conservative Semigroups via Generalized Doeblin’s Conditions. *Acta Appl. Math.*, 166:29–72, 2020.
- [9] V. Bansaye, J.-F. Delmas, L. Marsalle, and V. C. Tran. Limit theorems for Markov processes indexed by continuous time Galton-Watson trees. *Ann. Appl. Probab.*, 21(6):2263–2314, 2011.

- [10] J.-B. Bardet, A. Christen, A. Guillin, F. Malrieu, and P.-A. Zitt. Total variation estimates for the TCP process. *Electron. J. Probab.*, 18(10):1–21, 2013.
- [11] M. Benaim, B. Cloez, and F. Panloup. Stochastic approximation of quasi-stationary distributions on compact spaces and applications. *Ann. Appl. Probab.*, 28(4):2370–2416, 2018.
- [12] É. Bernard and P. Gabriel. Asymptotic behavior of the growth-fragmentation equation with bounded fragmentation rate. *J. Funct. Anal.*, 272(8):3455–3485, 2017.
- [13] É. Bernard and P. Gabriel. Asynchronous exponential growth of the growth-fragmentation equation with unbounded fragmentation rate. *J. Evol. Equ.*, 20(2):375–401, 2020.
- [14] J. Bertoin. On a Feynman-Kac approach to growth-fragmentation semigroups and their asymptotic behaviors. *J. Funct. Anal.*, 277(11):108270, 29, 2019.
- [15] J. Bertoin and A. R. Watson. A probabilistic approach to spectral analysis of growth-fragmentation equations. *J. Funct. Anal.*, 274(8):2163–2204, 2018.
- [16] G. Birkhoff. Extensions of Jentzsch’s theorem. *Trans. Amer. Math. Soc.*, 85:219–227, 1957.
- [17] F. Bouguet. A probabilistic look at conservative growth-fragmentation equations. In *Séminaire de Probabilités XLIX*, volume 2215, pages 57–74. Springer, Cham, 2018.
- [18] J. Broda, A. Grigo, and N. P. Petrov. Convergence rates for semistochastic processes. *Discrete Contin. Dyn. Syst. Ser. B*, 24(1), 2019.
- [19] M. J. Cáceres, J. A. Cañizo, and S. Mischler. Rate of convergence to an asymptotic profile for the self-similar fragmentation and growth-fragmentation equations. *J. Math. Pures Appl.*, 96(4):334 – 362, 2011.
- [20] D. Chafaï, F. Malrieu, and K. Paroux. On the long time behavior of the TCP window size process. *Stochastic Process. Appl.*, 120(8):1518–1534, 2010.
- [21] N. Champagnat and D. Villemonais. Exponential convergence to quasi-stationary distribution and  $Q$ -process. *Probab. Theory Related Fields*, 164(1-2):243–283, 2016.
- [22] N. Champagnat and D. Villemonais. General criteria for the study of quasi-stationarity. Preprint, arXiv:1712.08092, 2018.

- [23] B. Cloez. Limit theorems for some branching measure-valued processes. *Adv. in Appl. Probab.*, 49(2):549–580, 2017.
- [24] B. Cloez and P. Gabriel. On an irreducibility type condition for the ergodicity of nonconservative semigroups. *C. R. Math. Acad. Sci. Paris*, 358(6):733–742, 2020.
- [25] P. Collet, S. Martinez, S. Méléard, and J. San Martín. Stochastic models for a chemostat and long-time behavior. *Adv. Appl. Probab.*, 45(3):822–836, 2013.
- [26] P. Collet, S. Martínez, and J. San Martín. *Quasi-stationary distributions*. Probability and its Applications (New York). Springer, Heidelberg, 2013.
- [27] R. Dautray and J.-L. Lions. *Mathematical analysis and numerical methods for science and technology. Vol. 3*. Springer-Verlag, Berlin, 1990.
- [28] T. Debiec, M. Doumic, P. Gwiazda, and E. Wiedemann. Relative entropy method for measure solutions of the growth-fragmentation equation. *SIAM J. Math. Anal.*, 50(6):5811–5824, 2018.
- [29] P. Del Moral. *Feynman-Kac formulae*. Probability and its Applications (New York). Springer-Verlag, New York, 2004.
- [30] P. Del Moral. *Mean field simulation for Monte Carlo integration*, volume 126 of *Monographs on Statistics and Applied Probability*. CRC Press, Boca Raton, FL, 2013.
- [31] P. Del Moral and L. Miclo. On convergence of chains with occupational self-interactions. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 460(2041):325–346, 2004.
- [32] O. Diekmann, H. J. A. M. Heijmans, and H. R. Thieme. On the stability of the cell size distribution. *J. Math. Biol.*, 19(2):227–248, 1984.
- [33] W. Doblin. Éléments d’une théorie générale des chaînes simples constantes de Markoff. *Ann. École Norm. (3)*, 57:61–111, 1940.
- [34] J. L. Doob. Conditional brownian motion and the boundary limits of harmonic functions. *Bull. Soc. Math. France*, 85(4):431–458, 1957.
- [35] J. L. Doob. *Classical potential theory and its probabilistic counterpart: Advanced problems*, volume 262. Springer Science & Business Media, 2012.

- [36] R. Douc, E. Moulines, P. Priouret, and P. Soulier. *Markov chains*. Springer Series in Operations Research and Financial Engineering. Springer, Cham, 2018.
- [37] V. Dumas, F. Guillemin, and P. Robert. Limit results for Markovian models of TCP. In *GLOBECOM'01. IEEE Global Telecommunications Conference (Cat. No. 01CH37270)*, volume 3, pages 1806–1810. IEEE, 2001.
- [38] G. Dumont and P. Gabriel. The mean-field equation of a leaky integrate-and-fire neural network: measure solutions and steady states. *Nonlinearity*, 33:6381–6420, 2020.
- [39] J. Engländer, S. C. Harris, and A. E. Kyprianou. Strong law of large numbers for branching diffusions. *Ann. Inst. Henri Poincaré Probab. Stat.*, 46(1):279–298, 2010.
- [40] H. Engler, J. Prüss, and G. F. Webb. Analysis of a model for the dynamics of prions. II. *J. Math. Anal. Appl.*, 324(1):98–117, 2006.
- [41] G. Ferré. *Large Deviations Theory in Statistical Physics: Some Theoretical and Numerical Aspects*. PhD thesis, Université Paris-Est, 2019.
- [42] G. Ferré, M. Rousset, and G. Stoltz. More on the long time stability of Feynman-Kac semigroups. *Stoch. PDE: Anal. Comp.*, 2020.
- [43] G. Frobenius. Über Matrizen aus nicht negativen Elementen. *Berl. Ber.*, 1912:456–477, 1912.
- [44] P. Gabriel. Measure solutions to the conservative renewal equation. *ESAIM Proc. Surveys*, 62:68–78, 2018.
- [45] P. Gabriel and H. Martin. Periodic asymptotic dynamics of the measure solutions to an equal mitosis equation. Preprint arXiv:1909.08276, 2019.
- [46] P. Groisman and M. Jonckheere. Front propagation and quasi-stationary distributions for one-dimensional Lévy processes. *Electron. Commun. Probab.*, 23:11 pp., 2018.
- [47] M. Hairer. *Convergence of Markov Processes*. 2016. <http://www.hairer.org/notes/Convergence.pdf>.
- [48] M. Hairer and J. C. Mattingly. Yet another look at Harris’ ergodic theorem for Markov chains. In *Seminar on Stochastic Analysis, Random Fields and Applications VI*, volume 63 of *Progr. Probab.*, pages 109–117. Birkhäuser/Springer Basel AG, Basel, 2011.



- [49] A. J. Hall and G. C. Wake. A functional-differential equation arising in modelling of cell growth. *J. Austral. Math. Soc. Ser. B*, 30(4):424–435, 1989.
- [50] T. E. Harris. The existence of stationary measures for certain Markov processes. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Contributions to Probability Theory*, pages 113–124, Berkeley, Calif., 1956. University of California Press.
- [51] T. E. Harris. *The theory of branching processes*. Die Grundlehren der Mathematischen Wissenschaften, Bd. 119. Springer-Verlag, Berlin; Prentice-Hall, Inc., Englewood Cliffs, N.J., 1963.
- [52] A. G. Hart, S. Martínez, and J. San Martín. The  $\lambda$ -classification of continuous-time birth-and-death processes. *Adv. Appl. Probab.*, 35(4):1111–1130, 2003.
- [53] W. Huisinga, S. Meyn, and C. Schütte. Phase transitions and metastability in Markovian and molecular systems. *Ann. Appl. Probab.*, 14(1):419–458, 2004.
- [54] I. Kontoyiannis and S. P. Meyn. Spectral theory and limit theorems for geometrically ergodic Markov processes. *Ann. Appl. Probab.*, 13(1):304–362, 2003.
- [55] I. Kontoyiannis and S. P. Meyn. Large deviations asymptotics and the spectral theory of multiplicatively regular Markov processes. *Electron. J. Probab.*, 10:no. 3, 61–123, 2005.
- [56] I. Kontoyiannis and S. P. Meyn. Geometric ergodicity and the spectral gap of non-reversible Markov chains. *Probab. Theory Related Fields*, 154(1-2):327–339, 2012.
- [57] M. G. Kreĭn and M. A. Rutman. Linear operators leaving invariant a cone in a Banach space. *Amer. Math. Soc. Translation*, 1950(26):128, 1950.
- [58] T. Kurtz, R. Lyons, R. Pemantle, and Y. Peres. *A conceptual proof of the Kesten-Stigum theorem for multi-type branching processes*. Classical and Modern Branching Processes. Springer, New York, 1997.
- [59] P. Laurençot and B. Perthame. Exponential decay for the growth-fragmentation/cell-division equation. *Comm. Math. Sci.*, 7(2):503–510, 2009.
- [60] P. Maillard. Speed and fluctuations of n-particle branching brownian motion with spatial selection. *Probab. Theory Related Fields*, 166(3):1061–1173, 2016.
- [61] A. Marguet. A law of large numbers for branching Markov processes by the ergodicity of ancestral lineages. *ESAIM Probab. Stat.*, 23:638–661, 2019.

- [62] A. Marguet. Uniform sampling in a structured branching population. *Bernoulli*, 25(4A):2649–2695, 2019.
- [63] N. Marić. Fleming–viot particle system driven by a random walk on  $\mathbb{N}$ . *J. Stat. Phys.*, 160(3):548–560, 2015.
- [64] J. A. J. Metz and O. Diekmann. *The Dynamics of Physiologically Structured Populations*, volume 68 of *Lecture notes in Biomathematics*. Springer, 1st edition, August 1986.
- [65] S. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Cambridge University Press, Cambridge, second edition, 2009.
- [66] S. P. Meyn and R. L. Tweedie. Stability of Markovian processes. III. Foster-Lyapunov criteria for continuous-time processes. *Adv. in Appl. Probab.*, 25(3):518–548, 1993.
- [67] P. Michel, S. Mischler, and B. Perthame. General relative entropy inequality: an illustration on growth models. *J. Math. Pures Appl.*, 84(9):1235–1260, 2005.
- [68] S. Mischler and J. Scher. Spectral analysis of semigroups and growth-fragmentation equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(3):849–898, 2016.
- [69] E. Nummelin. *General irreducible Markov chains and nonnegative operators*. Cambridge University Press, Cambridge, 1984.
- [70] R. D. Nussbaum. Hilbert’s projective metric and iterated nonlinear maps. *Mem. Amer. Math. Soc.*, 75(391):iv+137, 1988.
- [71] O. Perron. Zur Theorie der Matrices. *Math. Ann.*, 64(2):248–263, 1907.
- [72] B. Perthame. *Transport equations in biology*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [73] B. Perthame and L. Ryzhik. Exponential decay for the fragmentation or cell-division equation. *J. Diff. Equations*, 210(1):155–177, 2005.
- [74] L. C. G. Rogers and D. Williams. *Diffusions, Markov processes and martingales: Volume 2, Itô calculus*, volume 2. Cambridge university press, 2000.
- [75] R. Rudnicki and M. Tyran-Kamińska. *Piecewise deterministic processes in biological models*. SpringerBriefs in Applied Sciences and Technology. Springer, Cham, 2017.

- [76] E. Seneta. *Non-negative matrices and Markov chains*. Springer Series in Statistics. Springer, New York, 2006. Revised reprint of the second (1981) edition.
- [77] E. A. van Doorn. Quasi-stationary distributions and convergence to quasi-stationarity of birth-death processes. *Adv. Appl. Probab.*, 23(4):683–700, 1991.
- [78] E. A. van Doorn. Birth–death processes and associated polynomials. *Journal of Computational and Applied Mathematics*, 153(1):497 – 506, 2003.
- [79] A. Velleret. Unique Quasi-Stationary Distribution, with a possibly stabilizing extinction. Preprint arXiv:1802.02409, 2018.
- [80] A. Velleret. Exponential quasi-ergodicity for processes with discontinuous trajectories. Preprint arXiv:1902.01441, 2019.
- [81] D. Villemonais. Minimal quasi-stationary distribution approximation for a birth and death process. *Electron. J. Probab.*, 20:18 pp., 2015.
- [82] G. F. Webb. An operator-theoretic formulation of asynchronous exponential growth. *Trans. Amer. Math. Soc.*, 303(2):751–763, 1987.
- [83] L. Wu. Essential spectral radius for Markov semigroups. I. Discrete time case. *Probab. Theory Related Fields*, 128(2):255–321, 2004.
- [84] A. A. Zaidi, B. Van Brunt, and G. C. Wake. Solutions to an advanced functional partial differential equation of the pantograph type. *Proc. A.*, 471(2179):20140947, 15, 2015.
- [85] H. Zhang and Y. Zhu. Domain of attraction of the quasistationary distribution for birth-and-death processes. *J. Appl. Probab.*, 50(1):114–126, 03 2013.